

PRODUCT MONOGRAPH

PrEXJADE[®]
(deferasirox)

Dispersible Tablets for Oral Suspension
125 mg, 250 mg, or 500 mg

Iron chelating agent

Novartis Pharmaceuticals Canada Inc.
Dorval, Quebec, H9S 1A9

Date of Preparation:
October 18, 2006

Control No: 231555

Date of Revision:
January 2, 2020

EXJADE and JADENU are registered trademarks

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
DOSAGE FORMS, COMPOSITION AND PACKAGING	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	10
DRUG INTERACTIONS	19
DOSAGE AND ADMINISTRATION	21
OVERDOSAGE	26
ACTION AND CLINICAL PHARMACOLOGY	26
STORAGE AND STABILITY.....	28
DOSAGE FORMS, COMPOSITION AND PACKAGING	28
PART II: SCIENTIFIC INFORMATION	30
PHARMACEUTICAL INFORMATION.....	30
CLINICAL TRIALS.....	30
DETAILED PHARMACOLOGY	38
TOXICOLOGY	41
REFERENCES	44
PART III: CONSUMER INFORMATION.....	48

PrEXJADE[®]
(deferasirox)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
oral	Dispersible tablets for oral suspension 125 mg, 250 mg, or 500 mg	Lactose monohydrate <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

EXJADE[®] (deferasirox) is indicated in the management of chronic iron overload in patients with transfusion-dependent anemias aged 6 years or older.

EXJADE[®] is also indicated in the management of chronic iron overload in patients with transfusion-dependent anemias aged two to five who cannot be adequately treated with deferoxamine.

EXJADE[®] is also indicated for the treatment of chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes aged 10 years and older.

Therapy with EXJADE should be initiated and maintained by physicians experienced in the treatment of chronic iron overload due to blood transfusions.

Pediatrics (2 to 16 years of age): There are limited data available on the use of EXJADE in children aged 2 to 5 (see **Special Populations – Pediatrics**). The overall exposure of EXJADE in young children (aged 2 to 5) was about 50% lower than in adults and this age group may require higher maintenance doses than are necessary in adults (see **Dosage and Administration**).

Geriatrics (≥ 65 years of age): Four hundred and thirty-one (431) patients ≥ 65 years of age have been studied in clinical trials of EXJADE (see **Special Populations – Geriatrics**). The pharmacokinetics of EXJADE have not been studied in elderly patients. In clinical trials, elderly patients experienced a higher frequency of adverse reactions than younger patients and should be monitored closely for adverse reactions that may require a dose adjustment.

CONTRAINDICATIONS

EXJADE is contraindicated in patients with estimated creatinine clearance <60 mL/min or serum creatinine >2 times the age-appropriate upper limit of normal.

EXJADE is contraindicated in high risk myelodysplastic syndrome (MDS) patients, any other MDS patient with a life expectancy < 1 year and patients with other hematological and non-hematological malignancies who are not expected to benefit from chelation therapy due to the rapid progression of their disease.

EXJADE is contraindicated in patients with platelet counts < 50 x 10⁹/L.

The use of EXJADE (deferasirox) is contraindicated in patients with hypersensitivity to the active substance, deferasirox, or to any of the excipients. For a complete listing of excipients, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the product monograph.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Therapy with EXJADE should be initiated and maintained by physicians experience in the treatment of chronic iron overload due to blood transfusions.

EXJADE is contraindicated in patients with moderate and severe renal impairment (see CONTRAINDICATIONS) and has not been studied in patients with severe hepatic impairment.

The following are clinically significant adverse events;

- Acute renal failure
- Hepatic failure
- Gastrointestinal haemorrhage and perforations

EXJADE dispersible tablets is a formulation of deferasirox with lower bioavailability compared to JADENU film-coated tablets. EXJADE requires a different dosing regimen and method of administration compared to JADENU. To avoid dosing errors, it is important that prescriptions of deferasirox specify both the type of formulation (dispersible tablet or film-coated tablet) and the prescribed dose in mg/kg/day.

General

The decision to remove accumulated iron should be individualized based on anticipated clinical benefits and risks of chelation therapy (see DOSAGE AND ADMINISTRATION).

The safety of EXJADE when administered with other iron chelation therapy has not been established.

No studies on the effects of EXJADE on the ability to drive or use machines have been performed. Patients experiencing dizziness should exercise caution when driving or operating machinery.

Carcinogenesis and Mutagenesis

See **TOXICOLOGY – Mutagenicity and Carcinogenicity sections.**

Cardiovascular

EXJADE has not been studied in patients with acute cardiac failure due to iron overload. Therefore, the use of EXJADE is not recommended in these patients.

Ear/Nose/Throat

Auditory disturbances (high-frequency hearing loss, decreased hearing) have been reported with EXJADE therapy (see **ADVERSE REACTIONS**). Auditory testing is recommended before the start of EXJADE treatment and thereafter at regular intervals.

Gastrointestinal

Gastrointestinal irritation may occur during EXJADE treatment. Upper gastrointestinal (GI) ulceration and haemorrhage and upper and lower GI perforations have been reported uncommonly in patients, including children and adolescents, receiving EXJADE. There have been rare reports of fatal GI haemorrhages and perforations. Fatal haemorrhages have been reported more frequently in elderly patients who had advanced hematologic malignancies and/or low platelet counts. Multiple ulcers have been observed in some patients and there have been reports of ulcers complicated with gastrointestinal perforation (see **ADVERSE REACTIONS**). Physicians and patients should remain alert for signs and symptoms of GI ulceration, perforation and haemorrhage during EXJADE therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected.

Caution should be exercised in patients who are taking EXJADE in combination with drugs that have known ulcerogenic potential, such as NSAIDs, corticosteroids, or oral bisphosphonates, and in patients receiving anticoagulants (see **DRUG INTERACTIONS**).

EXJADE contains lactose (1.1 mg lactose for each mg of deferasirox). This medicine is not recommended for patients with rare hereditary problems of galactose intolerance, severe lactase deficiency or glucose-galactose malabsorption.

Hematologic

There have been post-marketing reports (both spontaneous and from clinical trials) of cytopenias in patients treated with EXJADE. Most of these patients had preexisting hematologic disorders that are frequently associated with bone marrow failure (see **ADVERSE REACTIONS – Post-Market Adverse Drug Reactions**). The relationship of these episodes to treatment with

EXJADE is unknown. In line with the standard clinical management of such hematological disorders, complete blood count (CBC) should be available at baseline and blood counts should be monitored regularly during therapy. Dose interruption of treatment with EXJADE should be considered in patients who develop unexplained cytopenia. Reintroduction of therapy with EXJADE may be considered, once the cause of the cytopenia has been elucidated.

Hepatic/Biliary/Pancreatic

EXJADE is not recommended in patients with severe hepatic impairment (Child-Pugh C) (see **DOSAGE AND ADMINISTRATION – Dosing Considerations** and **ACTION AND CLINICAL PHARMACOLOGY – Special Populations and Conditions**). Elevations of serum transaminase levels (greater than 5 times the upper limit of normal) have been observed in 40 patients (6.1%; 40/652) receiving EXJADE in the context of 4 registration studies. In these patients, the transaminase levels were already $>5*ULN$ at baseline in 6 of the 40 patients. In 25 of the 40 patients, the transaminase levels at baseline were above the upper limit of normal but less than $5*ULN$.

Although uncommon (0.3%), elevations of transaminases greater than 10 times the upper limit of the normal range, suggestive of hepatitis, have been observed in clinical trials.

In a 5 year pediatric observational study, single events of elevations in ALT and AST suspected to be related to EXJADE were reported in 21.1% and 11.9%, respectively. Approximately 12% of patients on study required a dose reduction or interruption of EXJADE to manage the increase in transaminases and 2.7% of patients discontinued treatment.

There have been postmarketing reports of hepatic failure in patients treated with EXJADE. There are a total of 24 international reports of hepatic failure – 21 post-marketing reports and 3 reports from clinical studies. Two of the 24 cases were reported in Canada. Most reports of hepatic failure involved patients with significant comorbidities including liver cirrhosis and multi-organ failure; fatal outcomes were reported in some of these patients. As of the cut-off date above, no patient with normal baseline liver function or without additional life-threatening complications of their underlying disease has developed hepatic failure.

It is recommended that serum transaminases, bilirubin and alkaline phosphatase be monitored before the initiation of treatment, every 2 weeks during the first month and monthly thereafter. If there is an unexplained, persistent and progressive increase in serum transaminase levels, EXJADE treatment should be interrupted.

In the clinical trial and post-marketing settings, cases of serious acute pancreatitis were observed with and without documented underlying biliary conditions. A causal association to EXJADE could not be ruled out.

Immune

Rare cases of serious hypersensitivity reactions (such as anaphylaxis and angioedema) have been reported in patients receiving EXJADE, with the onset of the reaction occurring in the majority

of cases within the first month of treatment (see **ADVERSE REACTIONS – Post-Market Adverse Drug Reactions**). If hypersensitivity reactions occur, EXJADE should be discontinued and appropriate medical intervention instituted. EXJADE should not be reintroduced in patients who have experienced previous hypersensitivity reactions on deferasirox due to the risk of anaphylactic shock.

Ophthalmologic

Ocular disturbances (lens opacities, early cataracts, maculopathies) have been reported with EXJADE therapy (see **ADVERSE REACTIONS**). Ophthalmic testing (including fundoscopy) is recommended before the start of EXJADE treatment and thereafter at regular intervals.

Renal

EXJADE has not been studied in patients with renal impairment. EXJADE treatment has been initiated only in patients with serum creatinine within the age-appropriate normal range and therefore must be used with caution in patients with elevated serum creatinine levels (see **CONTRAINDICATIONS**).

EXJADE-treated patients experienced dose-dependent increases in serum creatinine. Increases in creatinine that were > 33% at ≥ 2 consecutive post baseline visits occurred at a greater frequency in EXJADE-treated patients compared to deferoxamine-treated patients (38% vs. 14%, respectively) in study 0107. In these beta-thalassemia patients, 94% of the creatinine elevations remained within the normal range. Under the dose adjustment instructions, dose reduction was required in one third of patients showing serum creatinine increase. In most patients undergoing dose reductions serum creatinine levels did not return to baseline; in 60% of patients undergoing dose reduction, serum creatinine remained elevated at > 33% without progression (see **ADVERSE REACTIONS - Abnormal Hematologic and Clinical Chemistry Findings**).

Cases of acute renal failure (some with fatal outcome) have been reported following the post-marketing use of EXJADE. There have been rare cases of acute renal failure requiring dialysis. For the fatal cases, it is impossible to completely exclude a contributory role of EXJADE to the renal impairment, although the fatalities in these critically ill patients could be attributable to other underlying diseases. The fact that there was an improvement after stopping the treatment in most of the cases with non-fatal acute renal failure is suggestive of a contributory role of EXJADE to these cases (see **ADVERSE REACTIONS – Post Market Adverse Drug Reactions**).

It is recommended that serum creatinine and/or creatinine clearance be assessed twice before initiating therapy. Weekly monitoring of serum creatinine and/or creatinine clearance is recommended in the first month after initiation or modification of therapy, and monthly thereafter. Patients with pre-existing renal conditions or patients who are receiving medicinal products that may depress renal function may be more at risk of complications. Care should be taken to maintain adequate hydration in patients (see **DOSAGE AND ADMINISTRATION – Dosing Considerations**). Dose reduction, interruption, or discontinuations should be considered

for elevations in serum creatinine (see **ADVERSE REACTIONS – Abnormal Hematologic and Clinical Chemistry Findings**).

Renal tubulopathy has been reported in patients treated with EXJADE. The majority of these patients were children and adolescents with beta-thalassemia and serum ferritin levels <1,500 microgram/L.

Tests for proteinuria should be performed monthly. As needed, additional markers of renal tubular function (e.g. glycosuria in non-diabetics and low levels of serum potassium, phosphate, magnesium or urate, phosphaturia, aminoaciduria) may also be monitored. Dose reduction or interruption may be considered if there are abnormalities in levels of tubular markers and/or if clinically indicated.

If there is a progressive increase in serum creatinine beyond the upper limit of normal, EXJADE should be interrupted (see **DOSAGE AND ADMINISTRATION**).

Skin

Serious skin reactions: Severe cutaneous adverse reactions (SCARs), including cases of Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and hypersensitivity vasculitis, as well as drug reaction with eosinophilia and systemic symptoms (DRESS) which could be life-threatening or fatal, and rare cases of erythema multiforme, have occurred during EXJADE treatment. Patients should be advised of the signs and symptoms of severe skin reactions, and be closely monitored. Upon suspicion of any SCAR, EXJADE should be discontinued immediately and should not be reintroduced.

Skin rashes: skin rashes may also appear during EXJADE treatment. For rashes of mild to moderate severity, EXJADE may be continued without dose adjustment, since the rash often resolves spontaneously. For more severe rash, where interruption of treatment may be necessary, EXJADE may be re-introduced after resolution of the rash, at a lower dose followed by gradual dose escalation.

Special Populations

Pregnant Women: There are no adequate and well-controlled studies conducted in pregnant women. No clinical data on exposed pregnancies are available for EXJADE. Studies in animals have shown some reproductive toxicity at maternally toxic doses (see **DETAILED PHARMACOLOGY – REPRODUCTION AND TERATOLOGY**). The potential risk for humans is unknown. It is therefore recommended that EXJADE should not be used during pregnancy. Patients taking oral contraceptives may be at risk of getting pregnant because EXJADE may decrease the efficacy of hormonal contraceptives (see **DRUG INTERACTIONS**).

Nursing Women: It is not known whether deferasirox is excreted in human milk. In an animal study, deferasirox and its metabolites were present in breast milk of rats following a 10 mg/kg oral dose. The concentration of deferasirox was approximately 20-fold higher in maternal milk

than in maternal plasma 4-8 hours post dose (see **DETAILED PHARMACOLOGY – REPRODUCTION AND TERATOLOGY**). Therefore, women should be advised against breast-feeding while taking EXJADE.

Pediatrics (2 to 16 years of age): There are limited data on the safety and effectiveness of EXJADE in pediatric patients aged 2 to 5 (see **CLINICAL TRIALS**). EXJADE has not been associated with growth retardation in children followed for up to 5 years in clinical studies. However, as a precautionary measure, body weight and longitudinal growth in pediatric patients should be monitored at regular intervals (every 12 months).

In a 5-year observational study in which 267 children aged 2 to <6 years (at enrollment) with transfusional hemosiderosis received deferasirox, there were no unexpected safety findings observed regarding adverse events or laboratory abnormalities with the exception of an increase in single events of elevated transaminases suspected to be related to EXJADE: 21.1% and 11.9% of pediatric patients had elevated alanine aminotransferase (ALT) and aspartate aminotransferase, respectively. Within the range of the known safety profile, increases in serum creatinine of >33% and above the upper limit of normal (ULN) on ≥ 2 consecutive occasions were observed in 3.1% of children and elevation of ALT greater than 5 times the ULN on ≥ 2 consecutive occasions was reported in 4.3% of children.

Geriatrics (≥ 65 years of age): Four hundred and thirty-one (431) patients ≥ 65 years of age have been studied in clinical trials of EXJADE. The majority of these patients had myelodysplastic syndrome (MDS, n= 393; β -thalassemia, n= 2; other anemias, n= 36). In general, caution should be used in elderly patients due to the greater frequency of decreased hepatic, renal, or cardiac function, concomitant disease or other drug therapy. In clinical trials, elderly patients experienced a higher frequency of adverse reactions than younger patients and should be monitored closely for adverse reactions that may require a dose adjustment.

Monitoring and Laboratory Tests

Serum Ferritin: Serum ferritin should be measured monthly to assess response to therapy and to evaluate for the possibility of overchelation of iron, although the correlation coefficient between serum ferritin and liver iron content (LIC) was 0.63, and changes in serum ferritin levels may not always reliably reflect changes in LIC. If the serum ferritin falls consistently below 500 microgram/L, temporary interruption of EXJADE therapy should be considered. Closer monitoring of serum ferritin levels as well as renal and hepatic function is recommended during periods of treatment with high doses and when serum ferritin levels are close to the target range. Dose reduction may be considered to avoid overchelation.

As with other iron chelator treatment, the risk of toxicity of EXJADE may be increased when inappropriately given to patients with a low iron burden or with serum ferritin levels that are only slightly elevated (see **DOSAGE AND ADMINISTRATION**).

Renal: It is recommended that serum creatinine be assessed twice before initiating therapy and monitored weekly for the first month followed by monthly thereafter. Tests for proteinuria should be performed monthly (see **Renal** section above).

Hepatic: It is recommended that serum transaminases, bilirubin and alkaline phosphatase be monitored before the initiation of treatment, every 2 weeks during the first month and monthly thereafter (see **Hepatic/Biliary/Pancreatic** section above).

Hematologic: In line with the standard clinical management of such hematological disorders, complete blood count (CBC) should be available at baseline and blood counts should be monitored regularly during therapy. Dose interruption of treatment with EXJADE should be considered in patients who develop unexplained cytopenia. Reintroduction of therapy with EXJADE may be considered, once the cause of the cytopenia has been elucidated (see **Hematologic** section above).

Auditory: Auditory disturbances (high-frequency hearing loss, decreased hearing) have been reported with EXJADE therapy (see **ADVERSE REACTIONS**).

Auditory testing is recommended before the start of EXJADE treatment and thereafter at regular intervals.

Ophthalmologic: Ocular disturbances (lens opacities, early cataracts, maculopathies) have been reported with EXJADE therapy (see **ADVERSE REACTIONS**). Ophthalmic testing (including fundoscopy) is recommended before the start of EXJADE treatment and thereafter at regular intervals.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Over 7000 patients have been treated with EXJADE (deferasirox) in clinical studies as of October 31, 2011. In the initial registration program, 652 patients were treated with EXJADE (deferasirox) in therapeutic studies lasting for a median of 366 days in pediatric and adult patients [52 patients between 2 and 5 years of age, 240 patients between 6 and 16 years of age, 330 patients between 17 to 65 years of age and 30 patients \geq 65 years]. These 652 patients included 421 with β -thalassemia, 99 with rare anemias, and 132 with sickle cell disease. Of these patients, 302 were male and 456 were Caucasian. In the sickle cell disease population, 89% of patients were black.

The most frequently occurring adverse events (all causalities) in the therapeutic trials of EXJADE were diarrhea, vomiting, nausea, headache, constipation, dyspepsia, abdominal pain, pyrexia, cough, proteinuria, increases in serum creatinine and transaminases, pruritis and skin rash. Gastrointestinal disorders, increases in serum creatinine and skin rash were dose related. Adverse events which most frequently led to dose interruption, dose adjustment, or discontinuation of therapy were skin rash, gastrointestinal disorders, infections, increased creatinine, and increased transaminases.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and

should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

In clinical trials in patients with transfusional iron overload, the most frequent reactions reported during chronic treatment with EXJADE in adult and pediatric patients include gastrointestinal disturbances in about 26% of patients (mainly nausea, vomiting, diarrhea, or abdominal pain), and skin rash in about 7% of patients. Mild, non-progressive, dose-dependent increases in serum creatinine occurred in 34% of patients (see **Abnormal Hematologic and Clinical Chemistry Findings**).

In clinical trials in patients with transfusional iron overload, elevations of liver transaminases as suspected drug-related adverse events were reported in about 2% of patients. The increases in liver transaminases were not dose-dependent. Forty percent of these patients had elevated levels (above the upper limit of normal) prior to receiving EXJADE. Elevations of transaminases greater than 10 times the upper limit of the normal range, suggestive of hepatitis, were uncommon (0.3%). High frequency hearing loss and lenticular opacities (early cataracts) have been observed in <1% of patients treated with EXJADE (see **WARNINGS AND PRECAUTIONS, Ear/Nose/Throat and Ophthalmologic**).

In a 1-year, randomized, double-blind, placebo-controlled study in patients with non-transfusion-dependent thalassemia syndromes, the most frequently reported AEs in the deferasirox 10 mg/kg/day group (at least 10%) were headache (16.4%), upper respiratory tract infection (14.5%), oropharyngeal pain (10.9%), pyrexia (10.9%), and rash (10.9%). Table 1 displays adverse events occurring in >5% of Exjade-treated patients.

Table 1 Adverse Events Occurring in >5% of Exjade-treated Patients in Study A2209

	EXJADE	EXJADE	Placebo	Placebo	Placebo
	5 mg/kg/day	10 mg/kg/day	5 mg/kg/day	10 mg/kg/day	Any dose
	N=55	N=55	N=28	N=28	N=56
	n (%)	n (%)	n (%)	n (%)	n (%)
Any AE	42 (76.4)	43 (78.2)	20 (71.4)	25 (89.3)	45 (80.4)
Headache	2 (3.6)	9 (16.4)	4 (14.3)	4 (14.3)	8 (14.3)
Upper respiratory tract infection	7 (12.7)	8 (14.5)	5 (17.9)	6 (21.4)	11 (19.6)
Oropharyngeal pain	4 (7.3)	6 (10.9)	0	2 (7.1)	2 (3.6)
Pyrexia	6 (10.9)	6 (10.9)	5 (17.9)	3 (10.7)	8 (14.3)
Rash	2 (3.6)	6 (10.9)	1 (3.6)	2 (7.1)	3 (5.4)
Diarrhoea	3 (5.5)	5 (9.1)	2 (7.1)	4 (14.3)	6 (10.7)
Fatigue	1 (1.8)	5 (9.1)	2 (7.1)	2 (7.1)	4 (7.1)
Nausea	4 (7.3)	5 (9.1)	1 (3.6)	6 (21.4)	7 (12.5)
Abdominal pain	2 (3.6)	4 (7.3)	1 (3.6)	3 (10.7)	4 (7.1)
Anaemia	3 (5.5)	4 (7.3)	0	2 (7.1)	2 (3.6)
Nasopharyngitis	5 (9.1)	4 (7.3)	2 (7.1)	3 (10.7)	5 (8.9)
Rhinitis	1 (1.8)	4 (7.3)	1 (3.6)	0	1 (1.8)
Abdominal pain upper	3 (5.5)	3 (5.5)	0	0	0
Dyspepsia	0	3 (5.5)	0	0	0
Gastroenteritis	1 (1.8)	3 (5.5)	0	2 (7.1)	2 (3.6)
Influenza	3 (5.5)	3 (5.5)	1 (3.6)	0	1 (1.8)
Insomnia	1 (1.8)	3 (5.5)	2 (7.1)	0	2 (3.6)

In Study 2209, one patient in the placebo 10 mg/kg group experienced an ALT increase to >5 x ULN and >2 x baseline (Table 2). Three Exjade-treated patients (all in the 10 mg/kg group) had 2 consecutive serum creatinine level increases >33% from baseline and >ULN. Serum creatinine returned to normal in all patients (in one spontaneously and in the other two after drug interruption).

Table 2 Number (%) of Patients with Increases in Serum Creatinine or SGPT/ALT in Study 2209

Laboratory Parameter	EXJADE 5 mg/kg (N=55) n (%)	EXJADE 10 mg/kg (N=55) n (%)	Placebo 5 mg/kg (N=28) n (%)	Placebo 10 mg/kg (N=28) n (%)	Placebo (N=56) n (%)
Serum Creatinine					
Creatinine increase (>33% from baseline and >ULN at ≥2 consecutive post-baseline values)	0 (0.0)	3 (5.5)	0 (0.0)	0 (0.0)	0 (0.0)
SGPT/ALT					
SGPT/ALT (>5 x ULN and >2 x baseline)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.6)	1 (1.8)

A total of 652 patients were treated with EXJADE (deferasirox) in therapeutic studies of adult and pediatric patients with β -thalassemia (n=421), rare anemias (n=99) and sickle cell disease (n=132). This population was 46% male, 70% Caucasian and included 292 patients ≤16 years of age. In the sickle cell disease population, 89% of patients were black. A total of 94% of β -thalassemia patients, 70% of patients with rare anemias, and 86% of patients with sickle cell disease patients received therapy for ≥ 48 weeks.

The data in Table 3 displays the adverse events, regardless of causality, occurring in >5% of patients in either treatment group in the primary efficacy study 0107 in which 296 β -thalassemia patients were treated with EXJADE and 290 patients received deferoxamine as an active comparator. Adverse events which most frequently led to dose interruption, dose adjustment, or discontinuation of therapy were skin rash, gastrointestinal disorders, infections, increased creatinine, and increased transaminases (see Abnormal Hematologic and Clinical Chemistry Findings). Discontinuations due to adverse events with a suspected relationship to EXJADE occurred in 7 patients.

Table 3 Adverse Events Occurring in >5% of β -thalassemia Patients in the Comparative Trial

Preferred Term	EXJADE N=296 n (%)	Deferoxamine N=290 n (%)
Pyrexia	56 (18.9)	69 (23.8)
Headache	47 (15.9)	59 (20.3)
Abdominal pain	41 (13.9)	28 (9.7)
Cough	41 (13.9)	55 (19.0)
Nasopharyngitis	39 (13.2)	42 (14.5)
Diarrhea	35 (11.8)	21 (7.2)
Creatinine increased ¹	33 (11.1)	0 (0)
Influenza	32 (10.8)	29 (10.0)

Nausea	31 (10.5)	14 (4.8)
Pharyngolaryngeal pain	31 (10.5)	43 (14.8)
Vomiting	30 (10.1)	28 (9.7)
Respiratory tract infection	28 (9.5)	23 (7.9)
Bronchitis	27 (9.1)	32 (11.0)
Rash	25 (8.4)	9 (3.1)
Abdominal pain upper	23 (7.8)	15 (5.2)
Pharyngitis	23 (7.8)	30 (10.3)
Arthralgia	22 (7.4)	14 (4.8)
Acute tonsillitis	19 (6.4)	15 (5.2)
Fatigue	18 (6.1)	14 (4.8)
Rhinitis	18 (6.1)	22 (7.6)
Back pain	17 (5.7)	32 (11.0)
Ear infection	16 (5.4)	7 (2.4)
Urticaria	11 (3.7)	17 (5.9)
¹ >33% increase compared to average baseline values		

The data in Table 4 displays the adverse events, regardless of causality, occurring in >1% in the pooled β -thalassemia patients by dose administered. The most frequently reported adverse events were abdominal pain, pyrexia and headache. In the 30 mg/kg dose group, the most frequently reported adverse events were abdominal pain, diarrhea and increased serum creatinine. Skin rash and ALT increase were the only adverse events that resulted in discontinuation.

Table 4 Most frequently reported AEs (>1% of all patients) – pooled β -thalassemia patients by dose administered

Preferred term	EXJADE 10 mg/kg N=143	EXJADE 20 mg/kg N=106	EXJADE 30 mg/kg N=172	All patients N=421
	Total	Total	Total	Total
	n (%)	n (%)	n (%)	n (%)
Abdominal pain	38 (26.6)	21 (19.8)	41 (23.8)	100 (23.8)
Pyrexia	47 (32.9)	31 (29.2)	20 (11.6)	98 (23.3)
Headache	37 (25.9)	20 (18.9)	26 (15.1)	83 (19.7)
Cough	38 (26.6)	17 (16.0)	25 (14.5)	80 (19.0)
Diarrhea	24 (16.8)	9 (8.5)	37 (21.5)	70 (16.6)
Nasopharyngitis	23 (16.1)	16 (15.1)	20 (11.6)	59 (14.0)
Vomiting	28 (19.6)	12 (11.3)	18 (10.5)	58 (13.8)
Rash	12 (8.4)	10 (9.4)	30 (17.4)	52 (12.4)
Nausea	11 (7.7)	11 (10.4)	28 (16.3)	50 (11.9)
Creatinine increased ¹	2 (1.4)	13 (12.3)	34 (19.8)	49 (11.6)
Laryngeal pain	20 (14.0)	12 (11.3)	17 (9.9)	49 (11.6)
Pharyngitis	28 (19.6)	9 (8.5)	10 (5.8)	47 (11.2)
Influenza	19 (13.3)	12 (11.3)	13 (7.6)	44 (10.5)
Rhinitis	28 (19.6)	8 (7.5)	6 (3.5)	42 (10.0)
URTI ²	9 (6.3)	8 (7.5)	24 (14.0)	41 (9.7)

Table 4 Most frequently reported AEs (>1% of all patients) – pooled β -thalassemia patients by dose administered

Preferred term	EXJADE 10 mg/kg N=143	EXJADE 20 mg/kg N=106	EXJADE 30 mg/kg N=172	All patients N=421
	Total	Total	Total	Total
	n (%)	n (%)	n (%)	n (%)
Bronchitis	7 (4.9)	9 (8.5)	20 (11.6)	36 (8.6)
Arthralgia	13 (9.1)	8 (7.5)	13 (7.6)	34 (8.1)
Back pain	9 (6.3)	16 (15.1)	9 (5.2)	34 (8.1)
Constipation	9 (6.3)	6 (5.7)	12 (7.0)	27 (6.4)
Fatigue	7 (4.9)	6 (5.7)	13 (7.6)	26 (6.2)
Ear infection	13 (9.1)	7 (6.6)	3 (1.7)	23 (5.5)
Tonsillitis	8 (5.6)	7 (6.6)	6 (3.5)	21 (5.0)
Post procedural pain	2 (1.4)	8 (7.5)	10 (5.8)	20 (4.8)
Acute tonsillitis	2 (1.4)	6 (5.7)	11 (6.4)	19 (4.5)
Asthenia	8 (5.6)	7 (6.6)	4 (2.3)	19 (4.5)
Gastroenteritis	8 (5.6)	6 (5.7)	5 (2.9)	19 (4.5)
Chest pain	2 (1.4)	8 (7.5)	8 (4.7)	18 (4.3)
Ear pain	3 (2.1)	5 (4.7)	4 (2.3)	12 (2.9)
Palpitations	1 (0.7)	4 (3.8)	7 (4.1)	12 (2.9)
Tachycardia	5 (3.5)	4 (3.8)	3 (1.7)	12 (2.9)
Transfusion reaction	7 (4.9)	3 (2.8)	2 (1.2)	12 (2.9)
Urticaria	3 (2.1)	4 (3.8)	5 (2.9)	12 (2.9)
Dyspepsia	4 (2.8)	3 (2.8)	4 (2.3)	11 (2.6)
Pain in extremity	5 (3.5)	3 (2.8)	3 (1.7)	11 (2.6)
Pruritus	3 (2.1)	4 (3.8)	4 (2.3)	11 (2.6)
Rhinorrhoea	1 (0.7)	6 (5.7)	3 (1.7)	10 (2.4)
Sinusitis	0	6 (5.7)	4 (2.3)	10 (2.4)
Transaminases increased	8 (5.6)	1 (0.9)	1 (0.6)	10 (2.4)
Urinary tract infection	2 (1.4)	1 (0.9)	7 (4.1)	10 (2.4)
Herpes simplex	3 (2.1)	1 (0.9)	5 (2.9)	9 (2.1)
Otitis media	2 (1.4)	1 (0.9)	6 (3.5)	9 (2.1)
Toothache	2 (1.4)	3 (2.8)	4 (2.3)	9 (2.1)
Anxiety	3 (2.1)	2 (1.9)	3 (1.7)	8 (1.9)
Bone pain	1 (0.7)	1 (0.9)	6 (3.5)	8 (1.9)
Conjunctivitis	6 (4.2)	1 (0.9)	1 (0.6)	8 (1.9)
Dyspnoea	0	2 (1.9)	6 (3.5)	8 (1.9)
Muscle cramp	1 (0.7)	0	7 (4.1)	8 (1.9)
Productive cough	4 (2.8)	3 (2.8)	1 (0.6)	8 (1.9)
Tooth abscess	2 (1.4)	0	6 (3.5)	8 (1.9)
Abdominal distension	1 (0.7)	0	6 (3.5)	7 (1.7)
Cholelithiasis	2 (1.4)	1 (0.9)	4 (2.3)	7 (1.7)
Enteritis	5 (3.5)	1 (0.9)	1 (0.6)	7 (1.7)
Epistaxis	4 (2.8)	1 (0.9)	2 (1.2)	7 (1.7)
Erythema	3 (2.1)	2 (1.9)	2 (1.2)	7 (1.7)
Hypoacusis	4 (2.8)	2 (1.9)	1 (0.6)	7 (1.7)
Insomnia	0	3 (2.8)	4 (2.3)	7 (1.7)
Vertigo	2 (1.4)	4 (3.8)	1 (0.6)	7 (1.7)

Table 4 Most frequently reported AEs (>1% of all patients) – pooled β -thalassaemia patients by dose administered

Preferred term	EXJADE 10 mg/kg N=143	EXJADE 20 mg/kg N=106	EXJADE 30 mg/kg N=172	All patients N=421
	Total	Total	Total	Total
	n (%)	n (%)	n (%)	n (%)
Alanine aminotransferase increased	4 (2.8)	2 (1.9)	0	6 (1.4)
Cardiac murmur	0	0	6 (3.5)	6 (1.4)
Depression	0	2 (1.9)	4 (2.3)	6 (1.4)
Dizziness	1 (0.7)	2 (1.9)	3 (1.7)	6 (1.4)
Dysmenorrhoea	0	3 (2.8)	3 (1.7)	6 (1.4)
Lymphadenopathy	2 (1.4)	1 (0.9)	3 (1.7)	6 (1.4)
Myalgia	1 (0.7)	1 (0.9)	4 (2.3)	6 (1.4)
Pharyngitis streptococcal	3 (2.1)	3 (2.8)	0	6 (1.4)
Proteinuria	1 (0.7)	1 (0.9)	4 (2.3)	6 (1.4)
Rash maculo-papular	0	3 (2.8)	3 (1.7)	6 (1.4)
Seasonal allergy	0	1 (0.9)	5 (2.9)	6 (1.4)
Abdominal discomfort	1 (0.7)	0	4 (2.3)	5 (1.2)
Contusion	2 (1.4)	0	3 (1.7)	5 (1.2)
Cystitis	1 (0.7)	1 (0.9)	3 (1.7)	5 (1.2)
Frequent bowel movements	1 (0.7)	1 (0.9)	3 (1.7)	5 (1.2)
Oedema peripheral	0	2 (1.9)	3 (1.7)	5 (1.2)
Respiratory tract infection	1 (0.7)	1 (0.9)	3 (1.7)	5 (1.2)
Syncope	2 (1.4)	2 (1.9)	1 (0.6)	5 (1.2)
Viral infection	1 (0.7)	1 (0.9)	3 (1.7)	5 (1.2)

¹ >33% increase compared to average baseline values

² Upper respiratory tract infection

Less Common Clinical Trial Adverse Drug Reactions (<1%)

The less common adverse events which occurred in clinical trials and considered to be related to EXJADE are listed below.

Cardiovascular: QT prolongation¹

General disorders: Pyrexia, oedema, fatigue

Ear and labyrinth disorders: Deafness

Eye disorders: Cataract, maculopathy, optic neuritis

Gastrointestinal: Duodenal ulcer, gastric ulcer (including multiple ulcers) gastritis, gastrointestinal haemorrhage, oesophagitis

Hepatic/Biliary/Pancreatic: Cholelithiasis, hepatitis, acute pancreatitis²

Nervous system: Dizziness

Psychiatric disorders: Anxiety, sleep disorder

Renal and urinary disorders: Renal tubular disorder (Fanconi syndrome)

Respiratory, thoracic and mediastinal disorders: Pharyngolaryngeal pain

Skin and subcutaneous tissue disorders: Pigmentation disorder, erythema multiforme

¹ Three cases of QT interval prolongation were reported in the clinical trials, however, a causal relationship to study drug was not established.

² Cases of serious acute pancreatitis were observed with and without documented underlying biliary conditions.

Abnormal Hematologic and Clinical Chemistry Findings

In the comparative study 0107, 113 patients treated with EXJADE had non-progressive increases in serum creatinine > 33% above baseline (Table 5). Twenty-five (25) patients required dose reductions. Increases in serum creatinine appeared to be dose-related. Of the 17 patients with elevations in SGPT/ALT levels > 5 times the ULN at consecutive visits, one discontinued EXJADE therapy. One patient experienced increases in transaminases to >10x ULN which normalized upon drug discontinuation but then increased sharply upon rechallenge. Increases in transaminases did not appear to be dose-related and most of these patients had elevated transaminases prior to receiving EXJADE therapy.

Table 5 Number (%) of patients with increases in SGPT/ALT or serum creatinine in study 0107

Laboratory parameter	EXJADE N=296 n (%)	Deferoxamine N=290 n (%)
Serum creatinine		
No. patients with creatinine > 33% and <ULN at ≥ 2 consecutive post-baseline visits	106 (35.8)	40 (13.8)
No. patients with creatinine increase > 33% and >ULN at ≥ 2 consecutive post-baseline visits	7 (2.4)	1 (0.3)
SGPT/ALT		
No. patients with SGPT/ALT >5 x ULN at ≥2 post-baseline visits	8 (2.7)	2 (0.7)
No. patients with SGPT/ALT >5 x ULN at ≥2 consecutive post-baseline visits	17 (5.7)	5 (1.7)

A total of 652 patients were treated with EXJADE (deferasirox) in clinical studies 107, 108, and 109. Of these patients, 237 (36%) had an increase in serum creatinine >33% on at least 2 consecutive visits, 68 (11%) of whom underwent dose reduction. The remainder returned to

serum creatinine <33% above baseline without dose reduction. Of the 68 patients who underwent dose reduction, 17 (25%) returned to normal, 41 (60%) remained elevated at >33% without progression and the remaining 10 (15%) fluctuated between baseline and 33%.

Based on limited data in patients with sickle cell disease (N=132) and other rare anemias (N=99), the type and frequency of adverse events observed were similar to those observed in patients with β -thalassemia. The adverse event profile in patients <16 years of age was similar to that seen in adults, regardless of disease state.

In 49 adult β -thalassemia patients treated for greater than 1 year and up to 3 years, the type and frequency of adverse events was similar to that seen in patients treated for up to 1 year.

Post-Market Adverse Drug Reactions

Cases of acute renal failure (some with fatal outcome) have been reported following the post-marketing use of EXJADE. Rarely biopsy proven interstitial nephritis has also been reported.

Hypocalcemia has been reported to occur during EXJADE therapy.

Spontaneously reported adverse reactions, presented below, are reported voluntarily and it is not always possible to reliably establish frequency or a causal relationship to drug exposure.

Post-Market Information

Since the International Birth Date (November 2, 2005), the cumulative exposure to marketed EXJADE is 123,619 patient-years as of October 31, 2011.

Paediatric population

Renal tubulopathy has been reported in patients treated with EXJADE. The majority of these patients were children and adolescents with beta-thalassemia and serum ferritin levels <1,500 microgram/L.

Renal and urinary disorders

Acute renal failure (mostly serum creatinine increases $\geq 2x$ upper limit of normal, and usually reversible after treatment interruption), hematuria, renal tubular necrosis.

Skin and subcutaneous tissue disorders

Stevens-Johnson syndrome, hypersensitivity vasculitis, urticaria, erythema multiforme, alopecia, toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS).

Immune system disorders

Hypersensitivity reactions (including anaphylaxis and angioedema)

Gastrointestinal disorders

Duodenal ulcer, gastric ulcer, gastrointestinal bleeding, gastrointestinal perforation,

Blood and lymphatic system disorders

Agranulocytosis, neutropenia, thrombocytopenia and aggravated anemia

Hepatic/Biliary/Pancreatic

Hepatic failure.

DRUG INTERACTIONS

Drug-Drug Interactions

Use with other iron chelator: The safety of Exjade when administered with other iron chelation therapy has not been established.

Use with Aluminum Containing Antacid Preparations: The concomitant administration of EXJADE and aluminum-containing antacid preparations has not been formally studied. Although deferasirox has a lower affinity for aluminum than for iron, EXJADE should not be taken with aluminum-containing antacid preparations (see **WARNINGS AND PRECAUTIONS**).

Use with Agents Metabolised through CYP3A4: In a healthy volunteer study, the concomitant administration of EXJADE and midazolam (a CYP3A4 substrate) resulted in a decrease of midazolam exposure by 17%. In the clinical setting, this effect may be more pronounced. Therefore, caution should be exercised when deferasirox is combined with substances metabolised through CYP3A4 (e.g. cyclosporine, simvastatin, hormonal contraceptive agents), due to a possible decrease in efficacy.

Use with Agents Inducing UDP-glucuronosyltransferase (UGT) Metabolism: In a healthy volunteer study, the concomitant administration of EXJADE (single dose of 30 mg/kg) and the potent UDP-glucuronosyltransferase (UGT) inducer rifampicin (repeated dose of 600 mg/day) resulted in a decrease of deferasirox exposure by 44% (90% CI: 37% - 51%). Therefore, the concomitant use of EXJADE with potent UGT inducers (e.g. rifampicin, phenytoin, phenobarbital, ritonavir) may result in a decrease in EXJADE efficacy. If EXJADE and a potent UGT inducer are used concomitantly, increases in the dose of EXJADE should be considered based on clinical response to therapy.

Use with Bile Acid Sequestrants: In a healthy volunteer study, the administration of cholestyramine after a single dose of deferasirox resulted in a 45% decrease in deferasirox exposure (AUC).

Use with Agents Metabolized by CYP2C8: In a healthy volunteer study, the concomitant administration of EXJADE (repeated dose of 30 mg/kg/day) and the CYP2C8 substrate repaglinide (single dose of 0.5 mg) resulted in an increase in repaglinide AUC and C_{max} by 131% and 62%, respectively. When EXJADE and repaglinide are used concomitantly, careful monitoring of glucose levels should be performed. An interaction between EXJADE and other CYP2C8 substrates like paclitaxel cannot be excluded.

Use with Agents Metabolized by CYP1A2: In a healthy volunteer study, the concomitant administration of EXJADE (repeated dose of 30 mg/kg/day) and the CYP1A2 substrate theophylline (single dose of 120 mg) resulted in an increase in theophylline AUC by 84% (90% CI: 73% to 95%). The single dose C_{max} was not affected, but an increase of theophylline C_{max} is expected to occur with chronic dosing. When EXJADE and theophylline are used concomitantly, monitoring of theophylline concentration and possible theophylline dose reduction should be considered. An interaction between EXJADE and other CYP1A2 substrates such as clozapine and tizanidine may be possible.

Use with busulfan: Based on Literature reports, concomitant administration of deferasirox and busulfan resulted in an increase of busulfan exposure (AUC). The AUC increase ranged approximately 40 to 150%. The mechanism of the interaction remains unclear. Caution should be exercised when deferasirox is combined with busulfan and the patient's plasma concentrations of busulfan should be monitored.

Use with Digoxin: In healthy volunteers, EXJADE had no effect on the pharmacokinetics of digoxin. The effect of digoxin on EXJADE pharmacokinetics has not been studied.

Use with Vitamin C: The concomitant administration of EXJADE and vitamin C has not been formally studied. Doses of vitamin C up to 200 mg were allowed in clinical studies without negative consequences. High doses of vitamin C should not be used.

Use with ulcerogenic potential drugs: Concomitant administration of EXJADE with drugs that have known ulcerogenic potential, such as NSAIDs, corticosteroids, or oral bisphosphonates, and use of EXJADE in patients receiving anticoagulants may increase the risk of gastrointestinal irritation (see **WARNINGS AND PRECAUTIONS**).

Use with hydroxyurea: The interaction of EXJADE with hydroxyurea has not been formally studied. No inhibition of deferasirox metabolism by hydroxyurea is expected based on the results of an *in vitro* study.

Drug-Food Interactions

EXJADE should be taken on an empty stomach at least 30 minutes before eating the first meal of the day, preferably at the same time each day.

EXJADE tablets for oral suspension can be dispersed in water, orange juice or apple juice. Dispersion of EXJADE in carbonated drinks or milk is not recommended due to foaming and slow dispersion, respectively.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions between EXJADE and gallium contrast media have not been studied. It is known that the results of gallium-67 imaging may be distorted by the iron chelator deferoxamine due to chelation of gallium-67. It is therefore recommended that EXJADE therapy be interrupted at least five days before gallium-67 scintigraphy.

DOSAGE AND ADMINISTRATION

EXJADE dispersible tablets for oral suspension and JADENU film-coated tablets are different formulations of deferasirox. EXJADE requires a different dosing regimen and method of administration compared to JADENU. If converting from JADENU to EXJADE dispersible tablets, see the **Dosing Considerations** section below.

To avoid dosing errors, it is important that prescriptions of deferasirox specify both the type of formulation (dispersible tablet or film-coated tablet) and the prescribed dose in mg/kg/day.

Recommended Dose and Dosage Adjustment

1. Transfusional iron overload

It is recommended that therapy with EXJADE (deferasirox) be started when a patient has evidence of chronic iron overload, such as the transfusion of approximately 100 mL/kg of packed red blood cells (approximately 20 units for a 40 kg patient) and a serum ferritin consistently >1000 µg/L. Doses should be in mg/kg and must be calculated and rounded to the nearest whole tablet size. EXJADE is available in three strengths (125, 250 and 500 mg).

The goals of iron chelation therapy are to remove the amount of iron administered in transfusions and, as required, to reduce the existing iron burden. The decision to remove accumulated iron should be individualized based on anticipated clinical benefit and risks of chelation therapy.

EXJADE should be taken on an empty stomach at least 30 minutes before eating the first meal of the day, preferably at the same time each day.

1.1 Starting Dose

The recommended initial daily dose of EXJADE is 10, 20, or 30 mg/kg/day body weight, depending on the patient's transfusion rate and the goal of treatment:

Patients requiring maintenance of an acceptable body iron level

- An initial daily dose of 10 mg/kg/day is recommended for patients receiving less than 7 mL/kg/month of packed red blood cells (approximately <2 units/month for an adult) and for whom the objective is maintenance of an acceptable body iron level.
- An initial daily dose of 20 mg/kg/day is recommended for patients receiving more than 7 mL/kg/month of packed red blood cells (approximately >2 units/month for an adult) and for whom the objective is maintenance of an acceptable body iron level.

Patients requiring reduction of iron overload

- An initial daily dose of 20 mg/kg/day is recommended for patients receiving less than 14 mL/kg/month of packed red blood cells (approximately <4 units/month for an adult) and for whom the objective is gradual reduction of iron overload.
- An initial daily dose of 30 mg/kg/day is recommended for patients receiving more than 14 mL/kg/month of packed red blood cells (approximately >4 units/month for an adult) and for whom the objective is gradual reduction of iron overload.

The dose dependent iron excretion (mg/kg/day) was calculated from the change in LIC over one year, the amount of blood transfused and the weight of the patient. Using two example patients of 20 kg and 50 kg, the amount of iron excreted over one year could be calculated in terms of mg/year and transfusion unit-equivalents/year (assuming that one unit of PRBC contains 200 mg iron). Thus in a 50 kg adult, doses of 10, 20 and 30 mg/kg for one year can remove the amount of iron contained in about 20, 36 and 55 units of blood, respectively (i.e. about 1.5, 3 and 4.5 units of blood per month, respectively). In a 20 kg pediatric patient, doses of 10, 20 and 30 mg/kg for one year can remove the amount of iron contained in about 8, 14 and 22 units of blood, respectively (i.e. about 0.6, 1.2 and 1.8 units of blood per month; or 6, 12 and 18 mL/kg/month, respectively).

Table 6 Study 0107: Iron excretion during one year (PP-2 population, biopsy)

Initial dose (mg/kg)	n	Iron excretion (mg/kg/day)	Iron excretion (mg/year)		Iron excretion (transfusion unit equivalents/year)	
			20kg patient	50 kg patient	20kg patient	50 kg patient
5	8	0.13 ± 0.10	939 ± 726	2349 ± 1816	4.7 ± 3.6	11.7 ± 9.1
10	44	0.22 ± 0.14	1572 ± 1055	3930 ± 2638	7.9 ± 5.3	19.6 ± 13.2
20	64	0.39 ± 0.15	2841 ± 1102	7102 ± 2756	14.2 ± 5.5	35.5 ± 13.8
30	108	0.60 ± 0.23	4378 ± 1712	10945 ± 4280	21.9 ± 8.6	54.7 ± 21.4

1.2 Dose Adjustment

It is recommended that serum ferritin be monitored every month and that the dose of EXJADE be adjusted if necessary every 3 to 6 months based on serum ferritin trends to minimize the risk of overchelation (see **WARNINGS AND PRECAUTIONS**). Dose adjustments should be made

in steps of 5 or 10 mg/kg and are to be tailored to the individual patient's response and therapeutic goals (maintenance or reduction of body iron burden). In patients with beta-thalassemia not adequately controlled with daily doses of 30 mg/kg, doses of up to 40 mg/kg may be considered.

If the serum ferritin falls consistently below 500 µg/L, consideration should be given to temporarily interrupting therapy with EXJADE. As with other iron chelator treatment, the risk of toxicity of EXJADE may be increased when inappropriately high doses are given in patients with a low iron burden or with serum ferritin levels that are only slightly elevated. Doses of EXJADE should not exceed 30 mg/kg per day since, with the exception of beta-thalassemia patients, there is limited experience with doses above this level (See **CLINICAL TRIALS**).

The Liver Iron Concentration (LIC) should be assessed periodically by an appropriate method such as biopsy or MRI in order to verify treatment response.

2. Non-transfusion-dependent thalassemia syndromes

Chelation therapy should only be initiated when there is evidence of iron overload (liver iron concentration (LIC) ≥ 5 mg Fe/g dry weight (dw) or serum ferritin consistently >800 microgram/L). In patients with no LIC assessment, caution should be taken during chelation therapy to minimize the risk of over-chelation. Doses should be in mg/kg and must be calculated and rounded to the nearest whole tablet size. EXJADE is available in three strengths (125, 250 and 500 mg).

EXJADE should be taken on an empty stomach at least 30 minutes before eating the first meal of the day, preferably at the same time each day.

2.1 Starting Dose

The recommended initial daily dose of EXJADE is 10 mg/kg body weight.

2.2 Dose Adjustment

It is recommended that serum ferritin be monitored every month to assess the patient's response to therapy and to minimize the risk of overchelation (see **WARNINGS AND PRECAUTIONS**). Every 3 to 6 months of treatment, consider a dose increase in increments of 5 to 10 mg/kg if the patient's LIC is ≥ 7 mg Fe/g dw, or serum ferritin is consistently $>2,000$ microgram/L and not showing a downward trend, and the patient is tolerating the drug well. The incidence of adverse effects increases with increasing dose. Experience with doses of 15mg/kg is limited. Doses above 20 mg/kg are not recommended because there is no experience with doses above this level in patients with non-transfusion-dependent thalassemia syndromes.

In patients in whom LIC was not assessed and serum ferritin is $\leq 2,000$ microgram/L, dosing should not exceed 10 mg/kg.

For patients in whom the dose was increased to >10 mg/kg, dose reduction is recommended to 10 mg/kg or less when LIC is <7 mg Fe/g dw or serum ferritin is ≤2,000 microgram/L.

Once a satisfactory body iron level has been achieved (LIC <3 mg Fe/g dw or serum ferritin <300 microgram/L), treatment should be interrupted. Treatment should be re-initiated when there is evidence from clinical monitoring that chronic iron overload is present.

3. Transfusional iron overload and non-transfusion-dependent thalassemia syndromes

Dosing Considerations

Conversion from JADENU to EXJADE: For patients who are currently on chelation therapy with JADENU tablets and converting to EXJADE deferasirox tablets (for oral suspension), the dose of EXJADE should be about 30% higher, rounded to the nearest whole tablet. The table below provides additional information on dosing conversion to EXJADE tablets.

Table 7 Dosing Conversion to EXJADE dispersible tablets

	JADENU Tablets (blue oval tablet)	EXJADE Tablets for oral suspension (white round tablet)
Transfusion-Dependent Iron Overload		
Starting Dose	14 mg/kg/day	20 mg/kg/day
Titration Increments	3.5–7 mg/kg	5–10 mg/kg
Maximum Dose	28 mg/kg/day	40 mg/kg/day
Non-Transfusion-Dependent Thalassemia Syndromes		
Starting Dose	7 mg/kg/day	10 mg/kg/day
Titration Increments	3.5–7 mg/kg	5–10 mg/kg
Maximum Dose	14 mg/kg/day	20 mg/kg/day

Geriatrics (≥ 65 years of age): The pharmacokinetics of EXJADE have not been studied in geriatric patients. The dosing recommendations for elderly patients are the same as described above. In clinical trials, elderly patients experienced a higher frequency of adverse reactions than younger patients and should be monitored closely for adverse reactions that may require a dose adjustment.

Pediatrics (2 to 16 years of age): The dosing recommendations for pediatric patients are the same as for adult patients. It is recommended that serum ferritin be monitored every month to assess the patient’s response to therapy and to minimize the risk of overchelation (see **WARNINGS AND PRECAUTIONS**). In children < 6 years of age, exposure was about 50% lower than adults. Since dosing is individually adjusted according to response this difference in exposure is not expected to have clinical consequences. Changes in weight of pediatric patients over time must also be taken into account when calculating the dose.

Patients with renal impairment: EXJADE has not been studied in patients with renal impairment (see **CONTRAINDICATIONS**). For adult patients, the daily dose of EXJADE should be reduced by 10 mg/kg if a non-progressive rise in serum creatinine by >33% above the average of the pre-treatment measurements is seen at two consecutive visits, and cannot be attributed to other causes. From those patients who underwent dose reduction, creatinine levels returned to baseline in only 25% of patients and in 60% of them, creatinine levels remained elevated >33% of the average pre-treatment levels. For pediatric patients, the dose should be reduced by 10 mg/kg if serum creatinine levels rise above the age-appropriate upper limit of normal at two consecutive visits. A total of 6 patients < 16 years developed creatinine levels >ULN during the core phase of the registration studies. Dose reductions were performed in 5 patients, in 4 of whom the levels returned to baseline. Creatinine levels fell to < ULN in the fifth patient but remained higher than baseline.

If there is a progressive increase in serum creatinine beyond the upper limit of normal, EXJADE therapy should be interrupted (see **ADVERSE REACTIONS – Abnormal Hematologic and Clinical Chemistry Findings**).

Patients with hepatic impairment EXJADE has been studied in a clinical trial in patients with hepatic impairment. For patients with moderate hepatic impairment (Child-Pugh B), the starting dose should be reduced by approximately 50%. EXJADE should not be used in patients with severe hepatic impairment (Child-Pugh C) (see **WARNINGS and PRECAUTIONS and ACTION AND CLINICAL PHARMACOLOGY – Special Populations and Conditions**). EXJADE treatment has been initiated only in patients with baseline liver transaminase levels up to 5 times the upper limit of normal range. The pharmacokinetics of deferasirox were not influenced by such transaminase levels. The treating physician should initiate treatment with a dose taking into account general dosing instructions together with the extent of hepatic impairment. Close monitoring of efficacy and safety parameters is recommended. It is recommended that serum transaminase, bilirubin and alkaline phosphatase be monitored before the initiation of treatment, every 2 weeks during the first month and monthly thereafter. If there is an unexplained, persistent, and progressive increase in serum transaminase levels EXJADE treatment should be interrupted.

Patients with skin rash: Skin rashes may occur during EXJADE treatment. Severe skin rashes may require interruption of EXJADE treatment.

Gender: Females have a moderately lower apparent clearance (by 17.5%) for deferasirox compared to males. Since dosing is individually adjusted according to response this difference in clearance is not expected to have clinical consequences.

Missed Dose

If a dose is missed it should be taken as soon as remembered on that day, and the next dose should be taken as planned. Doses should not be doubled to make up for a missed dose.

Administration

Reconstitution: EXJADE tablets should be completely dispersed by stirring in water, orange juice, or apple juice until a fine suspension is obtained. Doses of < 1 g should be dispersed in 100 mL of liquid and doses of > 1 g in 200 mL of liquid. After swallowing the suspension, any residue should be resuspended in a small volume of liquid and swallowed. Tablets must not be chewed, split, crushed or swallowed whole.

Incompatibilities: Dispersion in carbonated drinks or milk is not recommended due to foaming and slow dispersion, respectively.

OVERDOSAGE

In healthy volunteers, single doses of up to 40 mg/kg were tolerated.

Early signs of acute overdose are digestive effects such as abdominal pain, diarrhea, nausea and vomiting. Hepatic and renal disorders have been reported, including cases of liver enzyme and creatinine increased with recovery after treatment discontinuation. An erroneously administered single dose of 90 mg/kg led to Fanconi syndrome which resolved after treatment.

There is no specific antidote for deferasirox. Standard procedures for management of overdose (e.g. induction of emesis or gastric lavage) may be indicated as well as symptomatic treatment, as medically appropriate.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

EXJADE (deferasirox) is an orally active chelator that is highly selective for iron (as Fe³⁺). It is a tridentate ligand that binds iron with high affinity in a 2:1 ratio. Although its highest affinity is for iron, deferasirox has a significant affinity for aluminium. Deferasirox has very low affinity for zinc and copper, and there are variable decreases in the serum concentration of these trace metals after the administration of deferasirox. The clinical significance of these decreases is uncertain.

Pharmacodynamics

Pharmacodynamic effects tested in an iron balance metabolic study showed that deferasirox (10, 20 and 40 mg/kg/day) was able to induce net iron excretion (0.119, 0.329 and 0.445 mg Fe/kg body weight/d, respectively) within the clinically relevant range (0.1-0.5 mg Fe/kg/day). Iron excretion was predominantly fecal.

Daily treatment with EXJADE at doses of 20 and 30 mg/kg for one year in frequently transfused

adult and pediatric patients with beta-thalassemia led to reductions in indicators of total body iron; liver iron concentration was reduced by about 0.4 and 8.9 mg Fe/g liver (biopsy dry weight) on average, respectively, and serum ferritin was reduced by about 36 and 926 µg/L on average, respectively. At these same doses the ratios of iron excretion: iron intake were 1.02 (indicating net iron balance) and 1.67 (indicating net iron removal), respectively. EXJADE induced similar responses in iron-overloaded patients with other anemias. Daily doses of 10 mg/kg for one year could maintain liver iron and serum ferritin levels and induce net iron balance in patients receiving infrequent transfusions or exchange transfusions.

The effect of 20 and 40 mg/kg of deferasirox on QT interval was evaluated in a single-dose, double-blind, randomized, placebo-and active-controlled (moxifloxacin 400 mg), parallel group study in 182 healthy male and female volunteers aged 18-65 years. No evidence of prolongation of the QTc interval was observed in this study; however, the relevance of this study to long-term EXJADE use is unknown.

In patients with non-transfusion-dependent thalassemia syndromes and iron overload, treatment with EXJADE at a dose of 10 mg/kg/day for one year led to a reduction in mean liver iron concentration from baseline by -3.80 mg Fe/g dw, while an increase of 0.38 mg Fe/g dw was observed in patients treated with placebo. In addition, treatment with EXJADE at a dose of 10 mg/kg/day for one year led to a reduction in mean serum ferritin from baseline by -222.0 microgram/L, while an increase of 114.5 microgram/L was observed in patients treated with placebo.

In patients with cardiac iron deposition (MRI T2* <20 ms), treatment with EXJADE was shown to remove cardiac iron as demonstrated by progressive improvements in T2* values over 3 years of observation. In patients without cardiac deposition, EXJADE was shown to prevent clinically relevant cardiac iron deposition (maintenance of T2* at >20 ms) up to 1 year of observation, despite significant ongoing transfusion exposure.

Pharmacokinetics

Absorption: EXJADE (deferasirox) is absorbed following oral administration with a median time to maximum plasma concentration (t_{max}) of about 1.5 to 4 hours. The C_{max} and AUC of deferasirox increase approximately linearly with dose after both single administration and under steady-state conditions. Exposure to deferasirox increased by an accumulation factor of 1.3 to 2.3 after multiple doses. The absolute bioavailability (AUC) of deferasirox tablets for oral suspension is 70% compared to an intravenous dose.

Total exposure (AUC) was approximately doubled when taken along with a high-fat breakfast (fat content > 50% of calories) and increased by about 50% when taken along with a standard breakfast. The bioavailability (AUC) of deferasirox was moderately elevated when taken 30 minutes before meals with normal content (25% elevation) or high fat content (13% elevation). EXJADE must therefore be taken on an empty stomach at least 30 minutes before eating, preferably at the same time each day (see **DOSAGE and ADMINISTRATION**).

The total exposure (AUC) to deferasirox when taken after dispersion of tablets in orange juice or

apple juice was equivalent to the exposure after dispersion in water (relative AUC ratios of 103% and 90%, respectively).

Distribution: Deferasirox is highly (~99%) protein bound almost exclusively to serum albumin. The percentage of deferasirox confined to the blood cells was 5% in humans. The volume of distribution at steady state (V_{ss}) of deferasirox is 14.37 ± 2.69 L in adults.

Metabolism: Glucuronidation is the main metabolic pathway for deferasirox, with subsequent biliary excretion. Deconjugation of glucuronidates in the intestine and subsequent reabsorption (enterohepatic recycling) is likely to occur. Deferasirox is mainly glucuronidated by UGT1A1 and to a lesser extent UGT1A3. CYP450-catalysed (oxidative) metabolism of deferasirox appears to be minor in humans (about 8%). No evidence for induction or inhibition of CYP450 enzymes (CYP1A1, CYP1A2 and CYP2D6) at therapeutic doses has been observed. No inhibition of deferasirox metabolism by hydroxyurea was observed in an *in vitro* study. Deferasirox undergoes enterohepatic recycling.

Excretion: Deferasirox and metabolites are primarily (84% of the dose) excreted in the feces. Renal excretion of deferasirox and metabolites is minimal (8% of the dose). The mean elimination half-life ($t_{1/2}$) ranged from 8 to 16 hours.

Special Populations and Conditions

Hepatic Insufficiency: The average AUC of deferasirox in 6 subjects with mild hepatic impairment (Child-Pugh A) increased 16% over that found in 6 subjects with normal hepatic function, while the average AUC of deferasirox in 6 subjects with moderate hepatic impairment (Child-Pugh B) increased 76% over that found in 6 subjects with normal hepatic function. The average C_{max} of deferasirox in subjects with mild or moderate hepatic impairment increased 22% over that found in subjects with normal hepatic function (see **WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION – Dosing Considerations**). Efficacy of EXJADE was not studied in this pharmacokinetic investigation of subjects with hepatic impairment.

STORAGE AND STABILITY

Store at room temperature (15-30°C). Protect from moisture.

Keep in the original package. Keep in a safe place out of the reach of children and pets.

DOSAGE FORMS, COMPOSITION AND PACKAGING

EXJADE (deferasirox) dispersible tablets for oral suspension is available in three strengths: 125 mg, 250 mg and 500 mg.

125 mg:

Off-white, round, flat tablet with beveled edge and imprinted with “J 125” on one side and “NVR” on the other.

250 mg

Off-white, round, flat tablet with beveled edge and imprinted with “J 250” on one side and “NVR” on the other.

500 mg

Off-white, round, flat tablet with beveled edge and imprinted with “J 500” on one side and “NVR” on the other.


The inactive ingredients are: lactose monohydrate, crospovidone, povidone (K30), sodium lauryl sulphate, microcrystalline cellulose, colloidal silicon dioxide and magnesium stearate.

EXJADE (deferasirox) is supplied in blisters of 28 dispersible tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Common name:	Deferasirox
Chemical name:	4-[3,5-bis-(2-hydroxyphenyl)-[1,2,4]-triazol-1-yl]benzoic acid
Molecular formula:	C ₂₁ H ₁₅ N ₃ O ₄
Molecular mass:	373.37
Structural formula:	

Physicochemical properties:	white to slightly yellow powder
Solubility:	0.4 mg/mL at pH 7.40, 25°C
pH:	The pH of a 0.5% (m/V) suspension of deferasirox in water at 22-25°C is 4.10

CLINICAL TRIALS

Study demographics and trial design

Study 0107, was a 1-year, multi-centre, open-label, randomized, Phase III, active comparator control study to compare EXJADE (deferasirox) and deferoxamine in patients with β -thalassemia and transfusional hemosiderosis. Patients ≥ 2 years of age were randomized in a 1:1 ratio to receive either oral EXJADE at starting doses of 5, 10, 20 or 30 mg/kg once daily or subcutaneous DESFERAL[®](deferoxamine) at starting doses of 20 to 60 mg/kg for at least 5 days per week based on LIC (liver iron concentration) at baseline (2 to 3, >3 to 7, >7 to 14 and >14 mg Fe/g dry weight (dw)). Patients randomized to deferoxamine who had LIC values <7 mg Fe/g dw were permitted to continue on their prior deferoxamine dose, even though the dose may have been higher than specified in the protocol. Consequently, the ratio of EXJADE to

^{Pr}DESFERAL[®] is a registered trademark

deferoxamine doses for the two lower LIC categories was disproportionately low (1:4) compared to the two upper LIC categories (1:2). A total of 586 patients were randomized and treated (including 154 patients <16 years of age and received either EXJADE (296 patients) or deferoxamine (290 patients). There were no major differences in the baseline demographic characteristics between the groups. In both groups more than 97% of patients had received prior chelation therapy. Approximately two-thirds of each group was heavily iron overloaded as evidenced by an LIC value > 7 mg Fe/g dw at baseline.

Study 0108 was an open-label, non-comparative, phase II trial of efficacy and safety of EXJADE given for 1 year to patients with chronic anemias and transfusional hemosiderosis unable to be treated with deferoxamine. Similar to Study 0107, patients received 5, 10, 20, or 30 mg/kg per day of EXJADE based on baseline LIC. A total of 184 patients (adult and pediatric) were treated in this study: 85 patients with β -thalassemia and 99 patients with other congenital or acquired anemias (myelodysplastic syndromes, n=47; Diamond-Blackfan syndrome, n=30; other, n=22). Nineteen percent (N=35) of patients were <16 years of age (11 patients were ≥ 2 - < 6 years, 11 patients were 6 - < 12 years, and 13 patients were 12 - < 16 years) and 16% (N=30) of patients were ≥ 65 years of age. Thirty-seven patients had not received prior chelation therapy.

Study 0109 was a 1-year, open-label, randomized, Phase II, active comparator control study to compare EXJADE and deferoxamine in patients with sickle cell disease and transfusional hemosiderosis. As in Study 0107, patients received 5, 10, 20, or 30, mg/kg per day of EXJADE or subcutaneous deferoxamine at doses of 20 to 60 mg/kg for 5 days per week based on baseline LIC. The primary objective of this study was safety and tolerability of EXJADE (deferasirox) in this patient population. The population examined in study 0109 was adult and pediatric patients with sickle cell disease and chronic iron overload from repeated blood transfusions. This population included individuals receiving intermittent or regular transfusions. A total of 195 patients were randomized to receive either EXJADE (132 patients) or deferoxamine (63 patients) with the following distribution by age group: 7 patients were 2-< 6 years; 45 patients were 6 - < 12 years; 46 patients were 12 - <16 years; 96 patients were ≥ 16 years. There were no major differences in the patient populations randomized to receive either deferasirox or deferoxamine with regard to baseline demographics and disease characteristics. In both groups about 60% of patients had received prior chelation therapy. A somewhat higher percentage of deferasirox patients were heavily iron overloaded (LIC value > 7 mg Fe/g dw) at baseline when compared with deferoxamine (deferasirox 64%; deferoxamine 49%).

Relevant demographic characteristics for these studies are shown in **Table 8** and **Table 9**.

Table 8 Summary of patient demographics for clinical trials in chronic iron overload

Study #	Trial design	Dosage, route of administration and duration	Study patients (n=number)	Mean age \pm SD (Range)	Gender M/F
---------	--------------	--	---------------------------	---------------------------	------------

0107	open-label, randomized, Phase III, active comparator control study	EXJADE by baseline LIC: 5, 10, 20, or 30 mg/kg DFO by baseline LIC [‡] : 20-30, 25-35, 35-50, >50 mg/kg Duration: 52 weeks	EXJADE =296 DFO =290	17.2 ± 9.71 (2-53)	282 (48.1%) / 304 (51.9%)
0108	open-label, non-comparative, phase II trial of efficacy and safety	EXJADE by baseline LIC: 5, 10, 20, or 30 mg/kg Duration: 52 weeks	EXJADE =184	35.0 ± 22.4 (3-81)	93 (50.5%) / 91 (49.5%)
0109	open-label, randomized, Phase II, active comparator control study	EXJADE by baseline LIC: 5, 10, 20, or 30 mg/kg DFO by baseline LIC: 20-30, 25-35, 35-50, >50 mg/kg Duration: 52 weeks (ongoing)	EXJADE =132 DFO =63	19.2 ± 10.9 (3-54)	80 (41.0%) / 115 (59.0%)

[‡]LIC: liver iron concentration

SD: standard deviation

EXJADE: deferasirox

DFO: deferoxamine

Table 9 Number and % of patients treated with EXJADE by study (n=652)

EXJADE patients	Study 106 N = 40	Study 107 N = 296	Study 108 N = 184	Study 109 N = 132	All patients N = 652
Patients < 16 years	36 (90%)	154 (52%)	35 (19%)	67 (51%)	292 (45%)
Age group					
≥ 2 - < 6 years	7 (17.5%)	30 (10.1%)	11 (6.0%)	4 (3.0%)	52 (8.0%)
6 - < 12 years	13 (32.5%)	67 (22.6%)	11 (6.0%)	30 (22.7%)	121 (18.6%)
12 - < 16 years	16 (40.0%)	57 (19.3%)	13 (7.1%)	33 (25.0%)	119 (18.3%)
16 - < 50 years	4 (10.0%)	142 (48.0%)	99 (53.8%)	63 (47.7%)	308 (47.2%)
50 - < 65 years	0	0	20 (10.9%)	2 (1.5%)	22 (3.4%)
≥ 65 years	0	0	30 (16.3%)	0	30 (4.6%)

Study results

In the primary efficacy study 0107, treatment duration was 12 months. LIC, an accepted indicator of total body iron burden, was assessed at baseline and after 12 months of therapy by liver biopsy or non-invasively by biomagnetic susceptometry. Success rate, the primary efficacy endpoint, was defined as a reduction in LIC of ≥ 3 mg Fe/g dw for baseline values ≥ 10 mg Fe/g dw, reduction of baseline values between 7 and < 10 to < 7 mg Fe/g dw, or maintenance or reduction for baseline values <7 mg Fe/g dw. EXJADE was to be declared non-inferior to

deferoxamine if the lower limit of the 95% confidence interval (two-sided) of the difference in success rates was above -15%.

Table 10 Success rates for study 0107 (reduction or maintenance of LIC according to baseline and non-inferiority to deferoxamine)

	EXJADE	DFO
Biopsy & SQUID	n=276	n=277
Success rate (n (%))	146 (52.9)	184 (66.4)
95% CI	[47.0, 58.8]	[60.9, 72.0]
Difference and 95% CI	-13.5 [-21.6, -5.4]	
LIC < 7 mg Fe/g dw	n=85	n=87
Success rate (n (%))	34 (40.0)	72 (82.8)
95% CI	[29.6, 50.4]	[74.8, 90.7]
Difference [95% CI]	-42.8 [-55.9, -29.7]	
LIC ≥ 7 mg Fe/g dw	n=191	n=190
Success rate (n (%))	112 (58.6)	112 (58.9)
95% CI	[51.7, 65.6]	[52.0, 65.9]
Difference [95% CI]	-0.3 [-10.2, 9.6]	

EXJADE: deferasirox

DFO: deferoxamine

The primary efficacy population consisted of 553 patients (EXJADE n=276; deferoxamine n=277) who had LIC evaluated at baseline and 12 months or discontinued due to an AE. Of these 553 patients, 56 patients were < 6 years; 130 patients were 6 - < 12 years; 106 patients were 12 - <16 years; 261 patients were ≥ 16 years and <65 years. The overall success rates were 52.9% for EXJADE and 66.4% for deferoxamine with a difference of -13.5 in success rates and a 95% CI of [-21.6, -5.4]. Non-inferiority to deferoxamine was not achieved because the lower limit of the CI was below -15%. However, non-inferiority was demonstrated in a group of patients with baseline LIC levels ≥ 7 mg Fe/g dw who were allocated to the higher dose groups (EXJADE doses of 20 or 30 mg/kg and deferoxamine doses of ≥ 35 mg/kg. The success rates with EXJADE and deferoxamine were 58.6% and 58.9%, respectively, and the lower limit of the 95% CI (-10.2%) was above the non-inferiority threshold of -15% (see **Table 10**).

In patients with LIC ≥ 7 mg Fe/g dw who were treated with EXJADE 20 to 30 mg/kg per day a statistically significant reduction in LIC from baseline was observed (-5.3 ± 8.0 mg Fe/g dw, $p < 0.001$, t-test) which was not statistically significantly different from deferoxamine (-4.3 ± 5.8 mg Fe/g dw, $p = 0.367$).

Table 11 Ratio of iron excretion/iron intake and change in serum ferritin levels from baseline to 1 year of treatment in the primary efficacy study 0107


Protocol Recommended Dose (mg/kg/day)		Mean actual prescribed dose (mg/kg/day)		Ratio of iron excretion/iron intake		Serum ferritin levels (µg/L) Mean change from baseline ± SD	
EXJADE	Deferoxamine	EXJADE	Deferoxamine	EXJADE	Deferoxamine	EXJADE	Deferoxamine
				Mean ±	Mean ± SD	Mean ±	Mean ± SD

				SD (n)	(n)	SD (n)	(n)
5	20-30	6.2 ± 1.6	33.9 ± 9.9	0.58 ± 0.328 (15)	0.95 ± 0.101 (13)	+1189 ± 700 (15)	+211 ± 459 (13)
10	25-35	10.2 ± 1.2	36.7 ± 9.2	0.67 ± 0.365 (68)	0.98 ± 0.217 (75)	+833 ± 817 (73)	+32 ± 585 (77)
20	35-50	19.4 ± 1.7	42.4 ± 6.6	1.02 ± 0.398 (77)	1.13 ± 0.241 (87)	-36 ± 721 (80)	-364 ± 614 (89)
30	≥50	28.2 ± 3.5	51.6 ± 5.8	1.67 ± 0.716 (108)	1.44 ± 0.596 (98)	-926 ± 1416 (115)	-1003 ± 1428 (101)

SD: standard deviation

Reduction of LIC and serum ferritin were observed with EXJADE doses of 20 to 30 mg/kg. EXJADE doses below 20 mg/kg/day failed to provide consistent lowering of LIC and serum ferritin levels (Figure 1). Therefore, a starting dose of 20 mg/kg/day is recommended (see **DOSAGE AND ADMINISTRATION**).

Figure 1 Changes in Liver Iron Concentration and Serum Ferritin Following EXJADE (5 to 30 mg/kg per day) in Study 107



The results of the primary efficacy study are supported by the second major efficacy study, study 0108. The primary endpoint was to demonstrate a success rate significantly greater than 50% with EXJADE. In the total population, the success rate (50.5%) was not statistically significantly higher than 50%. However, in patients with LIC ≥ 7 mg Fe/g dw for whom both baseline and end of study LIC was available and who received EXJADE 20 to 30 mg/kg per day, the success rate was 58.5% [p=0.022 (50.3, 66.6)] and there was a statistically significant reduction in the absolute LIC from baseline to end of study (-5.5 ± 7.4 mg Fe/g dw, p < 0.001, t-test). There was

also a dose dependent effect on serum ferritin and the ratio of iron excretion to iron intake from doses of 5 to 30 mg/kg per day.

The primary objective of study 0109 was safety and tolerability (see **ADVERSE REACTIONS**). A total of 132 patients were treated with EXJADE and 63 patients with deferoxamine. At the time of the 6-month interim analysis, dose-dependent increases in the ratio of iron excretion to iron intake from doses of 5 to 30 mg/kg per day of EXJADE were observed. At the end of the study, the mean change in LIC in the per protocol-1 (PP-1) population, which consisted of patients who had at least one post-baseline LIC assessment, was -1.3 mg Fe/g dry weight for patients receiving EXJADE (n=113) and -0.7 mg Fe/g dry weight for patients receiving deferoxamine (n=54).

In an analysis of 192 beta-thalassemia patients dose escalated up to a maximum dose of 40 mg/kg/day (treated for up to 32 weeks), a further decrease in serum ferritin of 11.9% was observed (from the start of dosing >30 mg/kg/day). This was based on a pooled analysis of patients who were exposed to doses greater than 30 mg/kg/day in the key registration trials and their ongoing long-term extensions (Studies 0107/E, 0108/E, and 0109/E), and in another large clinical trial and its ongoing long-term extension (2402/E).

A cardiac sub-study was conducted as part of a Phase IV study. The cardiac sub-study was a one year, prospective, open-label, single-arm study which included two cohorts of severely iron overloaded β -thalassemia patients with LVEF values $\geq 56\%$: 114 patients with baseline T2* values >5 to <20 ms indicating myocardial siderosis (treatment cohort) and 78 patients with myocardial T2* ≥ 20 ms indicating no clinically significant cardiac iron deposition (prevention cohort). In the treatment cohort, the deferasirox starting dose was 30 mg/kg/day, with escalation to a maximum of 40 mg/kg/day. In the prevention cohort, the deferasirox starting dose was 20-30 mg/kg/day, with escalation to a maximum of 40 mg/kg/day. The primary endpoint of the cardiac sub-study was the change in T2* at one year. In the treatment cohort, T2* (geometric mean \pm coefficient of variation) significantly increased from a baseline value of 11.2 ms \pm 40.5% to 12.9 ms \pm 49.5%, representing a significant improvement of 16% ($p < 0.0001$). In the treatment cohort, improvement in T2* was observed in 69.5% of patients and stabilization of T2* in 14.3% of patients. LVEF remained stable and within the normal range: 67.4 \pm 5.7% to 67.1 \pm 6.0%. In the prevention cohort, myocardial T2* remained within the normal range and was unchanged from a baseline value of 32.0 ms \pm 25.6% to 32.5 ms \pm 25.1% (+2%; $p = 0.565$) indicating that daily treatment with deferasirox can prevent cardiac iron loading in β -thalassemia patients with a history of high transfusion exposure, and regular, ongoing transfusions.

Patients in the treatment cohort of the 1-year core study had the option to participate in two 1-year extensions. Over a three-year treatment duration period, there was a statistically significant ($p < 0.0001$), progressive improvement in the geometric mean of cardiac T2* from baseline overall, in the severe cardiac iron overload sub-group, which is associated with a high risk of cardiac failure (T2* >5 to <10 ms), and in the mild to moderate cardiac iron overload sub-group (T2* 10 to <20 ms) (Table 12). Using the geometric mean ratio, the T2* increase was 43% above baseline in all patients, 37% increase from baseline in the T2* >5 to <10 ms sub-group, and 46% increase from baseline in the T2* 10 to <20 ms sub-group. Continuous treatment with EXJADE for up to 3 years at doses >30 mg/kg/day effectively reduced cardiac iron in thalassemia major

patients with myocardial siderosis as shown by the number of patients who normalized their T2* or improved to a category associated with a lower risk of cardiac failure (Table 13).

Table 12 Geometric mean of T2* (ms) at baseline, and at the end of year 1, 2, and 3

Baseline cardiac T2* sub-group	Baseline (year 0)	End of core (year 1)	End of E1 (year 2)	End of E2 (year 3)
Overall	11.20 (n=105)	12.9(n=105) (p<0.0001)	14.79 (n=95) (p<0.0001)	17.12 (n=68) (p<0.0001)
T2* >5 to <10 ms	7.39 (n=41)	8.15 (n=41)	8.71 (n=35)	10.53 (n=24)
T2* 10 to <20 ms	14.62 (n=64)	17.39 (n=64)	20.13 (n=60)	22.32 (n=44)

E1 = end of first year extension

E2 = end of second year extension

Table 13 Transition table of cardiac T2* from core baseline to end of E2 (year 3)

Baseline cardiac T2* sub-group	Baseline n (%)	<5 ms n (%)	5 - <10 ms n (%)	10 - <20 ms n (%)	≥20 ms n (%)	Missing n (%)
>5 - <10 ms (N=39)	39 (100.0)	1 (2.6)	18 (46.2)	15 (38.5)	1 (2.6)	4 (10.3)
10 - <20 ms (N=62)	62 (100.0)		4 (6.5)	16 (25.8)	40 (64.5)	2 (3.2)
All patients (N=101)	101 (100.0)	1 (1.0)	22 (21.8)	31 (30.7)	41 (40.6)	6 (5.9)

A randomized, double-blind, placebo-controlled study to compare EXJADE and placebo was conducted in patients with non-transfusion-dependent thalassemia syndromes and iron overload. Patients ≥10 years of age were enrolled in the study in a 2:1:2:1 randomization to receive either EXJADE 5 mg/kg/day or EXJADE 10 mg/kg/day or matching placebo.

Transfusion independency of the patients was confirmed by the fact that blood transfusions 6 months prior to study start were not allowed and patients were excluded if a regular transfusion program was anticipated during the study. Iron overload was diagnosed by a serum ferritin >300 microgram/L at screening (two consecutive values at least 14 days apart from each other) and LIC ≥5 mg Fe/g dw measured by R2 MRI at screening. All patients with non-transfusion-dependent thalassemia syndromes were allowed with the exception of patients with HbS-variants or those whose clinical condition allowed phlebotomy.

In total, 166 patients were randomized. Demographics were well balanced. The main underlying disease was beta-thalassemia intermedia in 95 (57.2%) patients and HbE beta-thalassemia in 49 (29.5%) patients. The primary efficacy endpoint of change in liver iron concentration (LIC) from baseline to Week 52 was statistically significant in favor of both EXJADE treatment groups compared with placebo (Table 14). Furthermore, a statistically significant dose effect of EXJADE was observed in favor of the 10 mg/kg/day dose.

Table 14 Primary efficacy analysis – Analysis of covariance of absolute change in liver iron concentration (mg Fe/g dw) between baseline and Week 52 (Full Analysis Set)

	EXJADE 5 mg/kg/day (N=55)	EXJADE 10 mg/kg/day (N=55)	Placebo (N=56)
Change from baseline			
Number of evaluable patients	51	54	54
Least squares mean	-1.95	-3.80	0.38
Standard error	0.500	0.484	0.486
95% confidence interval	-2.94, -0.96	-4.76, -2.85	-0.59, 1.34
Difference of EXJADE - Placebo			
Least squares mean	-2.33	-4.18	-
Standard error	0.700	0.687	-
95% confidence interval (1)	-3.89, -0.76	-5.71, -2.64	-
p-value (2)	0.001	<.001	-
Difference of EXJADE 10 mg/kg – EXJADE 5 mg/kg			
Least squares mean	-	-1.85	-
Standard error	-	0.695	-
95% confidence interval	-	-3.22, -0.48	-
p-value (3)	-	0.009	-
<p>Estimates were obtained from an ANCOVA model for change in LIC between baseline and Week 52 with treatment as factor and baseline LIC as covariate.</p> <p>(1) two-sided simultaneous confidence intervals using Dunnett’s adjustment</p> <p>(2) one-sided p-value with Dunnett’s adjustment testing the hypothesis that the mean decrease in LIC is not greater under EXJADE than under placebo. Critical alpha-level: 0.025</p> <p>(3) two-sided p-value testing the hypothesis that the change in LIC is identical in the two EXJADE groups. Critical alpha-level: 0.05</p> <p>The last available post-baseline LIC was carried forward if no LIC value was available at Week 52.</p> <p>Only patients with both baseline and at least one post-baseline LIC value were included for this analysis.</p>			

The primary efficacy result was supported by additional analyses which showed a clear dose-response effect; this was reflected by a greater percentage of patients with an LIC decrease of ≥ 3 mg Fe/g dw in the 10 mg/kg/day EXJADE group compared to the 5 mg/kg/day EXJADE group (56.4% versus 32.7%, respectively). In addition, a reduction of $\geq 30\%$ in LIC between baseline and Week 52 was reported in approximately twice as many patients in the 10 mg/kg/day EXJADE group (49.15%) compared to the 5 mg/kg/day EXJADE group (25.5%). After one year of treatment, 27.3% of patients in the 10 mg/kg/day EXJADE group and 14.5% of patients in the 5 mg/kg/day EXJADE group achieved an LIC of < 5 mg Fe/g dw.

In the EXJADE treated groups, three pregnancies were reported among 45 female patients of child-bearing potential; one of these occurred despite concomitant oral contraceptive use. EXJADE may decrease the efficacy of hormonal contraceptives (see **DRUG INTERACTIONS**).

DETAILED PHARMACOLOGY

Pharmacology

EXJADE (deferasirox) is an orally active iron chelating agent. The core structure of deferasirox is an N-substituted bis-hydroxyphenyl-triazole, representative of a new class of tridentate and iron selective chelators. In this structure, potent iron-coordinating atoms are arranged in a geometry optimal for the formation of tridentate complexes.

Pharmacodynamics

In vitro

Affinity and selectivity of deferasirox for iron were assessed by potentiometric measurements, spectrophotometric titrations and cyclic voltammetry. Deferasirox has a high affinity for iron(III) with an overall affinity constant for the 1:2 complex (one Fe atom and two deferasirox molecules) in aqueous solution of 36.9 ($\log\beta_2$). Conversely, the affinity for iron(II) with a $\log\beta_2$ of 14.0 is low.

In a cell culture system using iron-loaded rat heart myocytes, deferasirox and deferoxamine showed similar potencies to remove iron at concentrations up to 80 $\mu\text{mol/L}$, which is a concentration that was achieved in human plasma following administration of efficacious doses.

In vivo

The potent and specific ability of deferasirox to mobilize tissue iron and to promote its excretion has been demonstrated in several animal studies. In the non iron-loaded, bile duct-cannulated rat, single oral doses of 25, 50 and 100 mg/kg deferasirox showed a rapid response within the first three hours after administration of the compound. A protracted action of biliary iron excretion was noted, extending beyond 24 hours for the high doses of 50 and 100 mg/kg. Furthermore, iron excretion was dose dependent. The efficiency of iron excretion, defined as the amount of iron excreted as a percentage of the theoretical iron binding capacity of the dose, was higher than previously tested compounds (deferoxamine s.c. 2-4% and oral L1 2%), and amounted to 18.3% for the 25 mg/kg dose, which showed the highest effect.

In iron-overloaded marmosets receiving 14, 28, 56 or 112 mg/kg deferasirox, significantly higher fecal iron was measured for the doses of 56 and 112 mg/kg even two days after administration of deferasirox, corroborating the prolonged action found in rats. In addition, a dose dependent increase of iron excretion and superior efficacy of deferasirox compared to other chelators was found. With both animal models the bulk of the iron was excreted into bile (rat) or feces

(marmoset) with less than 15% of the total iron found in urine, indicating that the iron complex is mainly cleared by the liver.

Radioactive iron given intravenously as deferasirox-iron complex was excreted in feces. By inference, this suggests that iron complexes formed with deferasirox in the blood are also cleared by the liver.

Chronic administration of deferasirox to rats and marmosets demonstrated effective removal of iron from the liver, the main storage organ for iron. Conversely, in marmosets, deferasirox did not reduce liver zinc or liver copper levels. Likewise, zinc and copper levels in kidney were not found to be negatively affected, whereas kidney iron levels were reduced by approximately 40% in males and 30% in females at the highest dose of 80 mg/kg tested.

Safety Pharmacology

In the course of its safety evaluation, it could be shown in rat that deferasirox does not promote the uptake of dietary iron. A wide range of safety pharmacology studies has been conducted to assess the effects of deferasirox on behavior, cardiovascular, renal, and respiratory systems.

In mice, deferasirox effects on CNS function included ataxia (≥ 100 mg/kg), slight head tremors (1000 mg/kg), and effects on step-through passive avoidance. *In vitro* receptor-binding assays showed that deferasirox at 10 $\mu\text{mol/L}$ only interacted weakly with kainate receptors and the channel site of NMDA receptors.

Renal evaluations in the rat after single doses up to 1000 mg/kg revealed no effects on the excretion of Cl^- , Na^+ and K^+ and urine volume. Intraduodenal administration of deferasirox at doses up to 1000 mg/kg to anesthetized rats demonstrated no effect on respiratory rate, tidal volume or minute volume. A variety of *in vitro* and *in vivo* studies were conducted to explore possible cardiovascular effects of deferasirox.

The data from the *in vitro* studies with isolated atria, heart or Purkinje fibers demonstrated no consistent pattern of changes. In an *in vivo* dog telemetry study, deferasirox demonstrated an increase in mean heart rate only at an exposure (C_{max}) of 734 $\mu\text{mol/L}$. No ECG changes were observed in marmoset toxicity studies after 4 weeks (130 mg/kg; C_{max} of 127-135 $\mu\text{mol/L}$) or 39 weeks (80 mg/kg; C_{max} of 64-81 $\mu\text{mol/L}$). Neither the hERG assay nor the dog study showed any evidence for QTc prolongation potential.

Pharmacokinetics

Pharmacokinetics and disposition of ^{14}C -labeled and non-radiolabeled deferasirox, its metabolites, and the respective iron complex $\text{Fe}[\text{deferasirox}]_2$ were investigated comprehensively in mice, rats, dogs and marmosets, including in humans. The fate of deferasirox appears similar in all species including human, with minor differences.

The extent of oral absorption and bioavailability of deferasirox was investigated after intravenous and oral administration of ¹⁴C-labeled deferasirox in mice, rats and marmosets and with nonradiolabeled deferasirox in dogs (see **Table 15**).

Table 15 Pharmacokinetic parameters of total deferasirox

	Human	Marmoset	Rat ^{b, f}
Single oral dose (mg/kg)	~ 20 ^a	25	10
AUC (μmol·h/L)/(mg/kg) ^c	35.5	8.72 ^h	2.71 ^h
C _{max} (μmol/L)/(mg/kg) ^c	1.53	1.85	0.76
t _{max} (h)	4-6	0.5	0.5
Bioavailability (% of dose)	73 ± 20 ^d	88 ^b	32 ^b
deferasirox(% of ¹⁴ C-AUC)	91	25	67
t _{1/2α} of deferasirox (h)	11 ± 5	0.7 (t _{1/2α} 35) ^{e, f}	0.8
	Human ^d	Marmoset	Rat ^{b, f}
Intravenous dose (mg/kg)	1.65	10	10
AUC (μmol·h /L)/(mg/kg) ^c	63.4	9.96 ^{b, k}	8.45 ^j
V _{ss} (L/kg)	0.18	8.1 ^{e, f}	0.64
Clearance CL (mL/min/kg)	0.74	3.2 ^f	5.6
Hepatic extraction ratio E _H (%) ^g	8	12	23
t _{1/2α} of deferasirox(h)	4.1 ± 1.5	0.5 (t _{1/2z} 51) ^{e, f}	0.7

^a: an oral dose of 1000 mg ¹⁴C-labeled deferasirox was given as a drink suspension in water to thalassemia patients at steady-state (daily 1000 mg non-radiolabeled deferasirox)

^b: calculations based on: deferasirox total = deferasirox free + Fe-[deferasirox]₂

^c: multiply μmol/L or μmol·h/L with 373.37 to obtain μg/L or μg·h/L, respectively

^d: healthy volunteers, 130 mg, 90 min i.v. infusion *versus* 375 mg, p.o.

^e: value very high, probably due to substantial contribution by enterohepatic recirculation evident in the terminal elimination phase after 8 h

^f: parameter calculated by the author

^g: E_H= CL/hepatic plasma flow, where hepatic plasma flow = hepatic blood flow . hematocrit (HCT ~0.45)

^h: AUC_{0-72h}; ⁱ AUC_{0-24h}; ^j AUC_{0.083-24h}; ^k AUC_{0.083-72h}

Using specific and sensitive analytical methods, deferasirox, metabolites and iron complex Fe-[deferasirox]₂ were quantified in various biological matrices. Orally administered deferasirox is well and rapidly absorbed in all species investigated including man. Oral bioavailability is substantial if not complete, with dose over-proportional increase in rodents and female rabbits, probably due to saturation of elimination processes. In marmosets and humans the systemic exposure to deferasirox increased proportionally to the dose. No unexpected accumulation and no significant gender differences were observed in the pharmacokinetics. Deferasirox is the major active circulating moiety in the animal species and human, and is considered to contribute most to the overall iron elimination *in vivo*. Two hydroxy metabolites of deferasirox (M1 and M2), which were found to be able to form iron complexes *in vitro* are considered to contribute negligibly to the overall iron excretion capacity of deferasirox.

Deferasirox in the blood was mainly confined to the plasma compartment of human and dog (≥ 90%), and to a lesser extent in the rabbit, marmoset, rat and mouse. For the deferasirox iron complex, almost no uptake to blood cells was observed. Deferasirox and its iron complex were extensively (98%-99%) bound to plasma proteins and primarily to human serum albumin for all species including human.

Deferasirox shows a distribution pattern typical for a compound with a low volume of distribution: deferasirox was distributed throughout the body, but was mainly present intravascularly. Substantial levels were found in organs of the gastrointestinal tract and excretory organs. Deferasirox and/or its metabolites passed the blood-brain barrier to a very low extent only. The placental barrier was passed to a very low extent only. Deferasirox was enriched in the milk building up a milk-to-plasma ratio of up to 20. Suckling juvenile rats were distinctly exposed to deferasirox. The tissue distribution pattern in juvenile animals was qualitatively similar to that in mother animals. No notable retention was observed in any tissue or organ of the albino and pigmented rat.

Metabolism of deferasirox includes mainly glucuronidation (animals and human), and to a lesser extent cytochrome P450-catalysed hydroxylation, in human mainly by CYP1A1, CYP1A2 and CYP2D6. Direct glucuronidation of deferasirox to the acyl-glucuronide (M3) occurred predominantly by UGT1A1 and UGT1A3. Drug-drug interactions by deferasirox based on UGT isoenzymes are in principle possible when a second co-administered drug is metabolized solely or mainly by UGT1A1 or UGT1A3. Any inhibition or induction of the cytochrome P450 enzymes by co-medications is not expected to significantly affect the pharmacokinetics of deferasirox. The potential for drug-drug interactions between deferasirox and comedications *via* cytochrome P450 enzymes, and *via* hepatic anion transport appears low. Based on the available data on the pharmacokinetic and disposition of deferasirox in animals and man, deferasirox appears to have a very low potential for induction of drug metabolizing enzymes in the liver.

Elimination of deferasirox and metabolites is rapid and complete. The elimination of the iron complex of deferasirox could not be determined in bile and/or feces due to its inherent instability in these matrices. Key elimination processes are hepatic metabolism and hepatobiliary elimination. Biliary elimination could be studied in rats only, but the findings are assumed to apply to higher animal species and humans as well. Hepatobiliary elimination may occur to some extent by first pass. There is evidence for enterohepatic recirculation of deferasirox and its metabolites. Enterohepatic recirculation can be ascribed to hepatobiliary elimination and intestinal hydrolysis of glucuronides to deferasirox. Deferasirox, its metabolites and the iron complex are anions, and seem to be eliminated largely *via* bile by hepatic canalicular anion transport (as shown in data from *mrp2*-deficient (TR-) rats). Active transporters expressed at the canalicular membranes of the hepatocytes *e.g.* MRP2, MXR (also called BCRP) may be involved in the elimination of deferasirox, its iron complex and its metabolites.

TOXICOLOGY

Acute Toxicity Studies

Single oral doses of deferasirox at 1000 mg/kg in mice and ≥ 500 mg/kg in rats resulted in mortality/morbidity. Single intravenous doses of deferasirox in mice resulted in mortality at 150 mg/kg. No mortality was observed in rats at the highest intravenous dose tested, 75 mg/kg.

Subacute Toxicity Studies

Mortality was observed at doses ≥ 200 mg/kg and at 100 mg/kg in the 2-week and 4-week rat study, respectively. Decreased tissue iron and changes in hematological parameters characteristic of a potent iron chelator were evident. Histopathologic findings of renal cortical tubular cytoplasmic vacuolation and gastrointestinal tract were common to both studies. Decreased hematopoiesis in the spleen, and splenic lymphoid depletion was observed after two weeks of administration. All effects were reversible following a nondosing phase. In a rat exploratory studies in which rats were iron overloaded or received diet supplemented with iron or findings were limited to pharmacological effects on tissue/serum iron levels.

In 2 and 4-week studies in marmosets, decreased tissue iron levels was observed at all doses of deferasirox. Effects on hematopoiesis were evident at 400 mg/kg after 2-weeks of administration and at 130 mg/kg after 4-weeks of treatment. vacuolar degeneration of the renal cortical tubules at doses ≥ 200 mg/kg and at 130 mg/kg in the 2-week and 4-week study, respectively. Vacuolation of intrahepatic bile duct cells and marked inflammation of gall bladder epithelium with fibrosis of the gall bladder wall and vacuolar hyperplasia of the epithelium was noted in a single animal at 130 mg/kg after 4 weeks treatment. All effects were reversible following a nondosing phase. In a two week exploratory study in marmosets preloaded with iron, no deferasirox related effects were observed. Dietary iron supplementation of marmosets did not reduce deferasirox effects.

Long Term Toxicity Studies

In a 26-week oral study in rats (with dietary iron supplementation) at doses of 0, 30, 80 or 180 mg/kg, mortality was observed at 180 mg/kg. Cataracts, characterized by lenticular degeneration and fragmentation, vacuole formation and/or lenticular epithelial hyperplasia were present at doses ≥ 80 mg/kg. Early lenticular changes were observed at 30 mg/kg. Cytoplasmic vacuolation of renal cortical tubular epithelium and splenic hematopoiesis occurred at 180 mg/kg. Ulceration/erosion of the glandular stomach was observed at ≥ 80 mg/kg. With the exception of the lenticular cataracts, all effects were reversible following a nondosing phase.

Oral administration of deferasirox to marmosets for 39 weeks at doses of 0, 20, 40 or 80 mg/kg resulted in mortality at 80 mg/kg. Histopathology findings at 80 mg/kg consisted of vacuolation of the hepatic bile duct cells; vacuolation and/or degeneration of the renal cortical tubules and dilatation of medullary tubules.

Fertility

Deferasirox at oral doses up to 75 mg/kg/day (which resulted in a drug exposure (plasma AUC) that was less than the maximum human value) was found to have no adverse effect on fertility and reproductive performance of male and female rats.

Reproduction and Teratology

Deferasirox was not teratogenic in rats or rabbits treated with doses up to and exceeding the maximum tolerated doses. Increased skeletal variations were seen in rats at a maternotoxic dose of 100 mg/kg/day, which achieved a drug exposure (plasma AUC) that was similar to the maximum human value. No adverse effect on fetal development was observed in rabbits at a maternotoxic dose of 50 mg/kg/day, which achieved a drug exposure about 30% of the maximum human value.

In a rat study designed to evaluate for effects on pre- and post-natal development, rats were treated at doses up to 90 mg/kg/day, a dose lethal to maternal animals, from early gestation to end of lactation. This treatment resulted in an increase in the number of stillborn pups and reduced pup birth weight.

Mutagenicity

Deferasirox was negative in the Ames test and an *in vitro* chromosome aberration assay with human peripheral blood lymphocytes. Positive responses were observed in an *in vitro* (V79) micronucleus screening test and in a rat *in vivo* bone marrow micronucleus assay, which may have been a result of altered hematopoiesis due to iron chelation. No response was observed in another rat *in vivo* micronucleus assay (liver) with doses up to 250 mg/kg.

Carcinogenicity

Deferasirox was not carcinogenic in a 104-week study in Wistar rats or in a 26-week study in transgenic p53^{+/-} heterozygous mice that were maintained on an iron-supplemented diet.

In the rat carcinogenicity study, rats were administered deferasirox daily for 2 years at doses up to 60 mg/kg resulting in plasma exposure that were 28 to 39% of human exposure at 20 mg/kg based on plasma AUC_{0-24hr}.

In the mouse oral carcinogenicity study, transgenic p53^{+/-} heterozygous mice were treated daily for 26 weeks at doses up to 200 mg/kg in males and 300 mg/kg in females, which resulted in plasma exposures that were 122% and 210% of human exposure at 20 mg/kg, respectively, based on plasma AUC_{0-24hr}.

104-week rat carcinogenicity study

No deferasirox-related neoplastic or non-neoplastic lesions were detected.

26-week transgenic mouse carcinogenicity study

No deferasirox-related neoplastic lesions were observed. Non-neoplastic lesions observed in mice were generally similar to those observed in 26 week toxicity study in rats and included biliary hyperplasia and hepatic periportal inflammation.

REFERENCES

1. [Adams RJ, McKie VC, Hsu L, et al (1998)] Prevention of a first stroke by transfusions in children with sickle cell anemia and abnormal results on transcranial Doppler ultrasonography. *N Engl J Med*, 339, 5-11.
2. [Aessopos A, Farmakis D, Hatziliami A, et al (2004)] Cardiac status in well-treated patients with thalassemia major. *Eur J Haematol*, 73, 359-66.
3. [Aldulak B, Bayazit AK, Noyan A, et al (2000)] Renal function in pediatric patients with β -thalassemia major. *Pediatr Nephrol*, 15, 109-112.
4. [Andrews NC (1999)] Disorders of iron metabolism. *N Engl J Med*, 341, 1986-95.
5. [Angelucci E, Brittenham GM, McLaren CE, et al (2000)] Hepatic iron concentration and total body iron stores in thalassemia major. *N Engl J Med*, 343, 327-31.
6. [Angelucci E, Brittenham GM, McLaren CE, et al (2000)] Hepatic iron concentration and total body iron stores in thalassemia major. *N Engl J Med* 343, 327-31. Erratum in: *N Engl J Med*, 2000, 343, 1740.
7. [Arboretti R, Tognoni G, Alberti D (2001)] Pharmacosurveillance and quality of care of thalassaemic patients. *Eur J Clin Pharmacol*, 56, 915-22.
8. [Ballas SK (2001)] Iron overload is a determinant of morbidity and mortality in adult patients with sickle cell disease. *Semin Hematol*, 38, 30-6.
9. [Borgna-Pignatti C, Rugolotto S, De Stefano P, et al (2004)] Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica*, 89, 1187-93.
10. [Brittenham GM, Griffith PM, Nienhuis AW, et al (1994)] Efficacy of deferoxamine in preventing complications of iron overload in patients with thalassemia major. *N Engl J Med*, 331, 567-73.
11. [Britton RS, Leicester KL, Bacon BR (2002)] Iron toxicity and chelation therapy. *Int J Hematol*, 76, 219-28.
12. [Bronspiegel-Weintrob N, Olivieri NF, Tyler B, et al (1990)] Effect of age at the start of iron chelation therapy on gonadal function in beta-thalassemia major. *N Engl J Med*, 323, 713-9.
13. [Cohen A (1987)] Management of iron overload in the pediatric patient. *Hematol Oncol Clin North Am*, 3, 521-44.

14. [Cohen AR, Galanello R, Piga A, et al (2000)] Safety profile of the oral iron chelator deferiprone: a multicentre study. *Br J Haematol*, 108, 305-12.
15. [Davis BA, Porter JB (2000)] Long-term outcome of continuous 24-hour deferoxamine infusion via indwelling intravenous catheters in high-risk beta-thalassemia. *Blood*, 95, 1229-36.
16. [Fischer R, Tiemann CD, Engelhardt R, et al (1999)] Assessment of iron stores in children with transfusion siderosis by biomagnetic liver susceptometry. *Am J Hematol*, 60, 289-99.
17. [Gabutti V, Piga A (1996)] Results of long-term iron-chelating therapy. *Acta Haematol*, 95, 26-36.
18. [Galanello R, Piga A, Alberti D, et al (2003)] Safety, tolerability, and pharmacokinetics of ICL670, a new orally active iron-chelating agent in patients with transfusion-dependent iron overload due to beta-thalassemia. *J Clin Pharmacol*, 43, 565-72.
19. [Harrison PM, Arosio P (1996)] The ferritins: molecular properties, iron storage function and cellular regulation. *Biochem Biophys Acta*, 1275, 161-203.
20. [Janka GE, Mohring P, Helmig M, et al (1981)] Intravenous and subcutaneous desferrioxamine therapy in children with severe iron overload. *Eur J Pediatr*, 137, 285-90.
21. [Jensen PD, Heickendorff L, Pedersen B, et al (1996)] The effect of iron chelation on haemopoiesis in MDS patients with transfusional iron overload. *Br J Haematol*, 94, 288-99.
22. [Koliakos G, Papachristou F, Koussi A, et al (2003)] Urine biochemical markers of early renal dysfunction are associated with iron overload in β -thalassemia. *Clin Lab Haem*, 25, 105-9.
23. [Kowdley KV (2004)] Iron, hemochromatosis, and hepatocellular carcinoma. *Gastroenterology*, 127, S79-86.
24. [Kwiatkowski JL, Cohen AR (2004)] Iron chelation therapy in sickle-cell disease and other transfusion-dependent anemias. *Hematol Oncol Clin North Am*, 18, 1355-77.
25. [Liu P, Olivieri N (1994)] Iron overload cardiomyopathies: new insights into an old disease. *Cardiovasc Drugs Ther*, 8, 101-10.
26. [National Comprehensive Cancer Network (2004)] Myelodysplastic syndromes. *Practice Guidelines in Oncology*. National Comprehensive Cancer Network, v1-2004.

27. [National Institutes of Health (2002)] Management and therapy of sickle cell disease. National Heart, Lung, And Blood Institute. NIH Publication No. 96-2117 (Fourth Edition).
28. [National Institutes of Health (2004)] NHLBI stops study testing how long children with sickle cell anemia should have blood transfusions to prevent stroke. NIH News, Clinical Alert, December 5, 2004.
29. [Nisbet-Brown E, Olivieri NF, Giardina PJ, et al (2003)] Effectiveness and safety of ICL670 in iron-loaded patients with thalassaemia: a randomized, double-blind, placebo-controlled, dose-escalation trial. *Lancet*, 361, 1597-1602.
30. [Olivieri NF (1999)] The β -thalassemias. *N Engl J Med*, 341, 99-109.
31. [Olivieri NF (2001)] Progression of iron overload in sickle cell disease. *Semin Hematol*, 38, 57-62.
32. [Olivieri NF and Brittenham GM. (1997)] Iron-chelating therapy and the treatment of thalassemia. *Blood*, 89:739-761.
33. [Olivieri, NF, Buncic JR, Chew E, et al (1986)] Visual and auditory neurotoxicity in patients receiving subcutaneous deferoxamine infusions. *N Engl J Med*, 314, 869-73.
34. [Olivieri NF, Nathan DG, MacMillan JH, et al (1994)] Survival in medically treated patients with homozygous beta-thalassemia. *N Engl J Med*, 331, 574-8.
35. [Pennell D, Porter B, Cappellini MD, et al] Deferasirox for up to 3 years leads to continued improvement of myocardial T2* in patients with beta-thalassemia major. *Haematologica*, haematol.2012 [Epub ahead of print]
36. [Piga A, Gaglioti C, Fogliacco E, Tricta F (2003)] Comparative effects of deferiprone and deferoxamine on survival and cardiac disease in patients with thalassemia major: a retrospective analysis. *Haematologica*; 88, 489-96.
37. [Porter J (2001)] Practical management of iron overload. *Br J Haematol*, 115, 239-52.
38. [Porter JB, Davis BA (2002)] Monitoring chelation therapy to achieve optimal outcome in the treatment of thalassaemia. *Best Pract Res Clin Haematol*, 15, 329-68.
39. [Schrier SL, Angelucci E (2005)] New strategies in the treatment of thalassemyias. *Annu Rev Med*, 56, 157-71.
40. [Sumboonnanonda A, Malasit P, Tanphaichitr VS, et al (1998)] Renal tubular function in β -thalassemia. *Pediatr Nephrol*, 12, 280-3.

41. [Taher AT, Porter J, Viprakasit V, et al (2012)] Deferasirox reduces iron overload significantly in nontransfusion-dependent thalassemia: 1-year results from a prospective, randomized, double-blind, placebo-controlled study. *Blood* 120, 970-977.
42. [Vichinsky E, et al. (2007)] A randomised comparison of deferasirox versus deferoxamine for the treatment of transfusional iron overload in sickle cell disease. *British Journal of Haematology*, 136, 501–508.

PART III: CONSUMER INFORMATION**PrEXJADE®**

(deferasirox dispersible tablets)

This leaflet is part III of a three-part "Product Monograph" published when **EXJADE®** was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about EXJADE. Contact your doctor or pharmacist if you have any questions about the drug.

Keep this leaflet. You may need to read it again. This medicine has been prescribed only for you or your child. Do not give it to anyone else or use it for any other illnesses.

ABOUT THIS MEDICATION**What the medication is used for:**

EXJADE is used to treat chronic iron overload in:

- adult patients and children aged 6 years and older who receive blood transfusions for the treatment of anemias;
- children aged 2 to 5 years who receive blood transfusions for the treatment of anemias, and who cannot be adequately treated with deferoxamine;
- adult patients and children aged 10 years and older with thalassemia syndromes who do not require regular blood transfusions for the treatment of anemia.

What it does:

EXJADE is an *iron chelating agent* which removes the excess iron from the body (also called iron overload), thereby reducing the risk of organ damage caused by iron overload.

When it should not be used:

- If you are allergic (hypersensitive) to deferasirox or any of the other ingredients (in particular, lactose) of EXJADE listed in the section *What the nonmedicinal ingredients are*.
- If you have severe kidney disease.
- If you have an advanced stage of myelodysplastic syndrome (MDS) or advanced cancer.
- If you have low platelet count ($<50 \times 10^9/L$).

What the medicinal ingredient is:

The active substance is deferasirox.

What the nonmedicinal ingredients are:

Lactose monohydrate, crospovidone, povidone, sodium lauryl sulphate, microcrystalline cellulose, silicon dioxide and magnesium stearate.

What dosage forms it comes in:

EXJADE is supplied as tablets for oral suspension. Each tablet contains 125 mg, 250 mg or 500 mg deferasirox.

Each blister package contains 28 dispersible tablets.

WARNINGS AND PRECAUTIONS**Serious Warnings and Precautions**

EXJADE should be prescribed by doctors experienced in the treatment of chronic iron overload due to blood transfusions.

EXJADE has not been studied in patients with severe kidney and liver problems (impairment).

Serious adverse events with the use of EXJADE include:

- acute kidney failure
- liver failure
- ulcer or bleeding in the stomach or intestines

BEFORE you use EXJADE talk to your doctor or pharmacist if you have:

- severe heart problems (acute cardiac failure).
- ulcer or bleeding in the stomach or intestines.
- liver or kidney problems.
- severe intolerance to lactose (milk sugars). EXJADE tablets contain lactose.
- visual (eye) problems.
- hearing problems.
- blood disorders (a low level of platelets or white blood cell count).
- skin problem.

During treatment with EXJADE, talk to your doctor or pharmacist immediately if you have:

- Rash, red skin, pain, swelling or blistering of the lips, eyes or mouth, skin peeling, high fever and flu-like symptoms and swollen lymph glands. If you get these symptoms, your doctor may stop your treatment.
- Vision and hearing problems.

Older people (65 years of age and above):

Elderly patients may experience more side effects than younger patients. They should be monitored closely by their doctor for side effects that may require a dose adjustment.

Children and adolescents (age 2 years to 16 years):

Their growth and development need to be monitored during treatment with EXJADE.

Pregnancy and breast-feeding:

EXJADE is not recommended during pregnancy unless clearly necessary. If you are pregnant or think that you may be, tell your doctor. EXJADE may decrease the effect of hormonal contraceptives, and you may be at risk of getting pregnant if you are taking a hormonal contraceptive. You should use an additional or different type of contraception.

Breast-feeding is not recommended during treatment with EXJADE.

Driving and using machines:

If you feel dizzy after taking EXJADE, do not drive or operate any tools or machines until you are feeling normal again.

You should receive regular blood and urine tests before and during treatment with EXJADE. You may also be assessed by Magnetic Resonance Imaging (MRI). These will monitor the amount of iron in your body (level of ferritin) to see how well EXJADE is working. The tests will also monitor your kidney function (blood level of creatinine, presence of protein in the urine) and liver function (blood level of transaminases, bilirubin and alkaline phosphatase). Your doctor will take these tests into consideration when deciding on the dose of EXJADE most suitable for you and will also use these tests to decide when you should stop taking EXJADE.

Your eyesight and hearing will also be tested before and periodically during treatment as a precautionary measure.

The safety of EXJADE when administered with other iron chelation therapy has not been established.

INTERACTIONS WITH THIS MEDICATION

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including non-prescription drugs (obtained without a prescription), vitamins and natural products. Some medicines may interact with EXJADE:

- Antacids (medicines used to treat heartburn) containing aluminum should not be taken at the same time of day as EXJADE.

In particular tell your doctor if you are taking any of the following:

- cyclosporine (used in transplantation to prevent graft rejection or for any other condition)
- simvastatin (used to lower cholesterol)
- hormonal contraceptive agents (birth control medicines)
- certain painkillers or anti-inflammatory medicines (e.g. acetylsalicylic acid, ibuprofen, corticosteroids)
- oral bisphosphonates (used to treat osteoporosis)
- anticoagulant medicines (used to prevent or treat blood clotting)
- repaglinide (used to treat diabetes)
- rifampicin (used to treat tuberculosis)
- paclitaxel (used in cancer treatment)
- phenytoin, phenobarbital (used to treat epilepsy)
- ritonavir (used in the treatment of HIV infection)
- cholestyramine (used mainly to lower cholesterol)
- theophylline (used to treat respiratory diseases such as asthma)
- busulfan (used as treatment prior to bone marrow transplant)

PROPER USE OF THIS MEDICATION

Always take EXJADE exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Usual dose:

For patients receiving regular blood transfusion:

- Initial dose: 10 mg, or 20 mg, or 30 mg per kg body weight daily.
- Maximum dose: 30 mg per kg body weight daily.

For patients with thalassemia syndromes who do not require regular blood transfusions:

- Initial dose: 10 mg per kg body weight daily.
- Maximum dose: 20 mg per kg body weight daily.

The daily dose will be adjusted depending on how you respond to the treatment.

When to take EXJADE

- Take EXJADE once a day, every day, at about the same time each day;
- Must be taken on an empty stomach;
- Then wait at least 30 minutes before eating the first meal of the day.

How to take EXJADE:

- **Drop** the tablet(s) into a glass of water, orange or apple juice (100 mL for doses of less than 1 g, and 200 mL for doses of 1 g or more).
- **Stir** until the tablet(s) dissolve completely. The liquid in the glass will look cloudy.
- **Drink** everything in the glass. Then add a little water or juice to what is left in the glass and drink that too.



Do not dissolve the tablets in fizzy drinks or milk.

Do not chew, break or crush the tablets. Do not swallow the tablets whole.

Overdose:

If you have taken too much EXJADE, or if someone else accidentally takes your tablets, contact your doctor or go to the hospital or contact your local poison control centre. Show the doctor the blister package of tablets. Urgent medical treatment may be necessary. You may experience effects such as abdominal pain, diarrhea,

nausea and vomiting, and kidney or liver problems that can be serious.

Missed Dose:

If you miss a dose, take it as soon as you remember on that day. Take your next dose as scheduled. Do not take a double dose on the next day to make up for the forgotten dose. Do not take more than one dose on the same day.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, EXJADE can cause side effects.

Common: *these side effects may affect between 1 and 10 in every 100 patients.*

- Gastrointestinal disorders, such as nausea, vomiting, diarrhea, pain in the abdomen, bloating, constipation, indigestion
- Skin rash
- Headache

Uncommon: *these side effects may affect less than 1 in every 100 patients.*

- Dizziness
- Fever
- Sore throat
- Swelling of arms or legs
- Change in the colour of the skin
- Anxiety
- Sleep disorder
- Tiredness
- Hearing loss
- Vision change (early cataracts)
- Ulcer and/or bleeding in the stomach or intestine
- Liver disorders
- Traces of blood and/or protein in the urine
- Hair loss
- A decrease in the number of cells involved in blood clotting (thrombocytopenia), in the number of red blood cells (anaemia aggravated), and in the number of white blood cells (neutropenia)

You will have some blood tests while taking EXJADE. Your doctor will look for any changes in kidney function, liver function, or in blood cell counts.

Your doctor may also want to test your eyesight and hearing while you are taking EXJADE.

You may notice other side effects not listed in this leaflet. If you are concerned with any side effect, or if any side effect makes you feel unwell, please tell your doctor or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Uncommon			
Blurred or cloudy eyesight		√	
Reduced hearing		√	
Severe upper stomach pain (sign of pancreatitis)		√	
Vomiting blood and/or have black stools		√	
Frequent heartburn or abdominal pain (signs of ulcers) particularly after eating or taking the drug		√	
Rare			
Acute renal failure (severe kidney problems), decreased urine output (sign of kidney problem)		√	
Difficulty breathing, dizziness, rash or swelling of the face and throat (signs of allergic reaction)			√
Frequent heartburn		√	
Partial loss of vision		√	
Rash, red skin, pain, blistering of the lips, eyes or mouth, skin peeling, high fever flu- like symptoms and swollen lymph glands (signs of serious skin reaction)			√
Very rare			
Drowsiness, upper right abdominal pain, yellowing or increased yellowing of your skin or eyes and dark urine (signs of liver problems)		√	
Unknown frequency			
Tear in stomach or intestine wall that can be painful and cause nausea			√

This is not a complete list of side effects. For any unexpected effects while taking EXJADE contact your doctor or pharmacist.

HOW TO STORE IT

- Keep out of the reach and sight of children and of

pets.

- Do not use EXJADE after the expiry date which is stated on the package/carton after EXP. The expiry date refers to the last day of that month.
- Store at room temperature (15-30°C).
- Store in the original package in order to protect from moisture.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhp-mps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at:

<http://www.novartis.ca>

or by contacting the sponsor, Novartis Pharmaceuticals Canada Inc., at: 1-800-363-8883

EXJADE and JADENU are registered trademarks

This leaflet was prepared by Novartis Pharmaceuticals Canada Inc. Dorval, Quebec, H9S 1A9

January 2, 2020