

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

P^rPATADAY[®]

olopatadine hydrochloride ophthalmic solution

Read this carefully before you start taking **PATADAY** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **PATADAY**.

What is PATADAY used for?

PATADAY is used for the treatment of itchy eyes caused by seasonal allergies.

How does PATADAY work?

Itchy eyes due to seasonal allergies, also called allergic conjunctivitis, happens when allergens like pollen cause the cells of the eye to release a chemical called histamine. This can result in itching, redness and swelling on the surface of your eye. PATADAY works by stopping the release of histamine and other chemicals that cause the allergic reaction. This reduces eye itching.

What are the ingredients in PATADAY?

Medicinal ingredients: olopatadine hydrochloride

Non-medicinal ingredients: benzalkonium chloride (preservative), disodium phosphate, edetate disodium, hydrochloric acid and/or sodium hydroxide (to adjust pH), povidone, purified water, sodium chloride.

PATADAY comes in the following dosage forms:

Ophthalmic solution (eye drops); 0.2% w/v

Do not use PATADAY if:

- you are allergic to olopatadine hydrochloride or to any of the other ingredients (see **What are the ingredients in PATADAY?**).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take PATADAY. Talk about any health conditions or problems you may have, including if you:

- are pregnant or planning to become pregnant.
- are breastfeeding. PATADAY may pass into breastmilk.
- are under 18 years of age.

Other warnings you should know about:

- **Use of PATADAY and use of contact lenses:**
 - Do not wear contact lenses if your eyes are red.
 - PATADAY contains a preservative, benzalkonium chloride, which may cause eye irritation and is known to discolour soft contact lenses. Do not use PATADAY while wearing contact lenses.
 - Remove your contacts before using PATADAY and wait at least 15 minutes before putting your contacts back in.
- **Use of PATADAY with other eye drops or ointments:**
 - If you use other eye drops, wait at least 5 minutes between putting in PATADAY and the other drops.
 - Apply eye ointments last.
- **Driving and using machines:** You may find that your vision is blurred for a time just after you use PATADAY. Do not drive or use machines until your vision is clear.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with PATADAY:


There are no known drugs that interact with PATADAY.

How to take PATADAY:


- PATADAY is an eye drop. Only use it in your eye(s).
- Use PATADAY exactly how your healthcare professional has told you to. Do not change your dose without talking to your healthcare professional.

Usual dose:

Adults: 1 drop in each affected eye once daily.



1



2

Instructions for use:

1. Get the PATADAY bottle and a mirror if needed.
2. Wash your hands.
3. Twist off the cap, being careful not to touch the dropper tip.
4. Hold the bottle, pointing it down, between your thumb and middle finger.
5. Tilt your head back. Pull down your eyelid with a clean finger, until there is a 'pocket' between the eyelid and your eye. The drop will go in there (Figure 1).
6. Bring the bottle tip close to the eye. Use the mirror if it helps.
Do not touch your eye or eyelid, or any surface with the dropper. It could contaminate the drops, cause an eye infection and damage the eyes.
7. Gently press the bottom of the bottle with your forefinger to release one drop (Figure 2). Do not squeeze the bottle: it is designed so that just a gentle press on the bottom is all that it needs.
8. If you use drops in both eyes, repeat the steps for the other eye.
9. Put the bottle cap firmly back on immediately after use.

Overdose:

If you get too much in your eyes, rinse it all out with warm water. Don't put in any more drops until it's time for your next regular dose.

If you think you, or a person you are caring for, have taken too much PATADAY, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to use PATADAY, use a single drop as soon as you remember, and then go back to your regular routine. **Do not** use a double dose to make up for the one missed.

What are possible side effects from using PATADAY?

These are not all the possible side effects you may have when taking PATADAY. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

In the eye:

- eye problems such as dry, itchy, red, irritated or crusted eyes
- eye surface inflammation with or without surface damage
- eye discharge
- eye pain
- increased tear production
- eyelid redness, swelling
- sensitivity to light

- blurred vision
- burning, stinging or gritty feeling or a feeling as if something is trapped in the eye

Other areas of your body:

- headache
- dizziness
- fatigue or tiredness
- nasal dryness
- a dry mouth
- a change in your sense of taste
- nausea
- red or itchy skin

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
Allergic reactions: swelling of the face, lips, tongue, mouth or throat, shortness of breath, difficulty swallowing, hives, severe itching and rash, nausea, vomiting			✓
Increased heart rate		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.